



Standard Heterosis for Grain Yield and its Attributing Traits in Early Maturing Maize Hybrids

Viqar-Un-Nisa ^a, Z. A. Dar ^a, S. Naseer ^a, M. H. Khan ^a, A. A. Lone ^a,
M. A. Wani ^a, M. A. Rather ^a, A. M. I. Qureshi ^a, Wajhat-Un-Nisa ^a,
Bilal Ahmad Lone ^{a*}, Aflaq Hamid ^a, Gowher Ali ^a, Asma Fayaz ^b
and S. Kumar ^a

^a Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir, India.
^b Chandigarh University, Chandigarh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2021/v33i2430796

Editor(s):

- (1) Prof. Rusu Teodor, University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca, Romania.
- (2) Prof. Hakan Sevik, Kastamonu University, Turkey.
- (3) Dr. Hon H. Ho, State University of New York, USA.

Reviewers:

- (1) Zziwa Simon, International Potato Center, Uganda.
- (2) Cousin Musvosvi, University of KwaZulu-Natal, South Africa.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:
<https://www.sdiarticle5.com/review-history/79633>

Original Research Article

Received 09 October 2021
Accepted 18 December 2021
Published 21 December 2021

ABSTRACT

The present study was conducted at the research farm of Dryland Agriculture Research Station, SKUAST-Kashmir during the kharif of 2019. Range of heterosis was calculated over standard check. The most desirable cross combinations viz., KDM-440 x KDM-914A (-5.879), KDM-440 x V-335 (-4.468), KDM-930 x V-351(-4.165), KDM-927A x V-335 (-3.986), CML-470 x KDM-914A (-3.808) for days to maturity, KDM-347 x V-351 (4.967) CML-470 x KDM-914A (4.610), CML-474 x V-351(-4.396), KDM-927 A x V-351 (-3.110), KDM-930 x V 335 (2.896) for number of kernel rows per cob, KDM-347 x V-351(-7.544), KDM-440 x V-351 (6.438), KDM-916A x KDM-914 A(6.117), CML-470 x V-351(5.992), CML-474 x V-335 (5.274) for number of kernels per row, KDM-340 x V-351 (3.966), KDM340 x KDM-914A (3.150), KDM-930 x V-351(3.165), KDM-440 x KDM-914 (3.556), KDM-347 x V-351(3.324) for 100-grain weight, KDM-347 x V351 (34.197), KDM-440 x V335 (28.933), CML470 x KDM-914A (29.380), KDM-895 x KDM914A (25.383), KDM927A x KDM-914A (19.549) for grain yield per plant were observed in the present study. The range of heterosis

over standard check for days to maturity from -3.796 (CML-425 × KDM-914A) to (2.414) CML-474 × KDM-914A. The extent of heterosis for number of kernel rows per ear over standard check ranged from CML-470 × V-335(28.576) to KDM-916A × V -335(-2.143), for quantity of kernels per row maximum well known heterosis ranged from CML-470 × V-351 (19.079) to KDM-347 × V-351 (-15.132), for 100- grain weight, heterosis ranged from 22.3% for CML-470 × KDM-914 A to -3.5% for KDM-916A × V-335. For grain yield per plant, heterosis ranged from 61.9% for CML-470 × KDM-914 A to 9.3% for KDM916A × V-335. Standard heterosis for crosses CML -470 × KDM-914 A, CML-474 × V-351, KDM-440 × V-335 should be tested for multilocation to make valid conclusion related to their use in commercial maize cultivation.

Keywords: Heterosis; hybrids; yield and yield attributes; early maturing; maize.

1. INTRODUCTION

Heterosis is the hybrid vigour which is generally represented as performance of offspring over parents. Charles Darwin had achieved lot of experiments in order to check his concept of origin of species. In which comparison of inbred lines with open pollinated maize was once compared. From this they concluded that cross pollinated vegetation has gained 25 percentage higher than offspring of self pollinated crops and in addition had super cold tolerance. Later in 1876 it was concluded from these experiments that hybrids outshined in discipline because of "great innate constitutional vigour" (Darwin et al., 1989). Meanwhile two very eminent scientists of time laid foundations for new maize breeding methods. Scientists describe hybrid vigour as it was very important to have some term for it so "Shull" was the first person who marked the "heterosis" term for the rationalization of hybrid vigour. "East and Shull" working independently in 1908 had rediscovered phenomena of inbreeding depression and hybrid vigour in maize, consequently hybrids as soon as identified would possibly be reproduced year after year except a change. In maize hybrids can be developed with the aid of de tasselling of one of the inbred and pollination is carried from different inbred lines planted in close by adjacent blocks.

Heterozygosis is accountable for diversity which makes the crosses to range from each other. Where as homozygosis is diminished mechanically due to inbreeding [1-4]. Generally F₁ hybrids of "outcrossed" inbred strains show a incredible increase in vigour and viability. Even if the inbred lines, are degenerated due to inbreeding and there will be decrease of vigour. This phenomenon is termed heterosis or hybrid vigour. So, for assessment in deciding the parents, crosses and any future breeding programme. With this aim and objective problem

was undertaken to assessemnet the heterosis for grain yield and its attributing traits in early maturing maize hybrids.

2. MATERIALS AND METHODS

Ten diverse maize inbred lines viz., KDM-340, KDM-440, KDM-916, KDM-927A, KDM-895, CML-470, CML-474, CML-425, KDM-347, KDM-930 and three widely adapted testers V-351, V-335, KDM-914A were crossed in line x tester mating design in order to obtain a set of thirty crosses during *rabi* season of 2018 at Winter Nursery, Agricultural centre of Research Institute, Rajendranagar, Hyderabad then F₁ was evaluated at Dryland Agriculture Research Station Srinagar during *khari* season of 2019 in randomized block design with three replications with spacing of 75 x 20 cm. The data was recorded on morphological, maturity, yield and yield attributing traits viz., days to 50 per cent tasseling, days to 50 per cent silking, days to maturity (Seed to seed), plant height (cm), ear height (cm): ear length (cm), ear diameter (cm), number of kernel rows ear⁻¹, number of kernels row⁻¹, prolificacy index: grain yield plant⁻¹ (g). Combinig ability procedure was done according to kemphrone (1957).

3. RESULTS AND DISCUSSION

Heterosis is regarded as the ideal performance of offspring over its parents and is the effect of variances. It is set up as a primary tool for enhacement of manufacturing of crops. Plant breeders have extensively used and exploited heterosis in analyzing yield of numerous plants in particular cross pollinated crop species. Therefore, the heterotic studies can supply the groundwork for the utilization of precious hybrid mixtures in the future breeding programmes and their profitable utilization. The presence of sufficient hybrid vigour is an full-size pre-requisite for triumphing manufacturing of hybrid varieties.

Table 1. Estimation of heterosis over standard check for days to maturity, yield & yield attributing traits in maize (*Zea mays* L.)

Crosses	Days to maturity	Kernel rows cob ⁻¹	Number of kernels row ⁻¹	100- grain weight(g)	Grain yield plant ⁻¹ (g)
CML470 × V-351	-3.793*	14.286*	19.079*	19.321*	51.154*
CML470 × KDM-914 A	1.690	15.195*	18.421*	22.321*	61.923*
CML470 × V335	0.992	28.576*	19.316*	6.071*	41.50*
CML474 × V351	-0.172*	10.714*	12.500*	14.821*	58.077*
CML474 × KDM-914A	-2.414*	3.571*	15.132*	10.250	50.769*
CML474Xv335	-3.241	7.143*	9.868*	13.143*	39.231*
KDM-916A xV335	-2.052*	-2.234	-5.921	-3.464	9.231
KDM-347x V351	-0.517*	9.700	-15.132	9.607*	13.462
KDM-347x KDM-914 A	-1.897*	-3.571	6.579*	12.214*	22.692*
KDM-347 xV335	2.241	5.809	5.921*	17.321*	48.462*
KDM-895 × V351	3.103	-3.670	4.605*	7.250*	33.846*
KDM-895 × KDM-914A	4.276	14.286*	11.842*	15.536*	45.769*
KDM-895 × V335	4.897	5.122*	-1.316	11.607*	18.077*
KDM-440 xV351	-1.962*	19.245*	15.789	6.964*	28.846*
KDM-440 × KDM914A	9.034	12.700	-10.526	11.443*	53.462*
KDM-440 xV335	1.823	8.571*	11.421*	9.571*	31.154*
KDM-930A × V351	-4.991*	6.243*	12.500*	12.500*	24.615*
KDM -930A × KDM914A	1.552	12.234*	5.550*	14.718*	37.308*
KDM930AxV335	-4.203*	7.183*	2.921*	10.107*	25.769*
KDM-340 × V351	2.241	2.193*	17.763*	14.821*	13.846
KDM-340 × KDM-914A	-0.142*	6.445*	24.652*	15.000*	36.215*
KDM-340 × V335	-0.163*	4.700	7.895*	18.893*	26.992*

In the cutting-edge reading, preferred heterosis viz., parental heterosis for each trait were computed. The consequences confirmed a wide array of patterns of heterosis for maturity the traits underneath study (Table 1). Maximum amount of heterosis was once calculated for general check for maturity traits. Out of thirty cross combinations, noteworthy heterosis over standard check (SMH 2) was recorded in KDM-440 × KDM-914A (-5.879), KDM-440 × V-335 (-4.468), KDM-930 × V-351(-4.165), KDM-927 × V-335 (-3.986), CML-470 × KDM-914A (-3.808) for Days to maturity. KDM-347 × V-351 (4.967), CML-470 × KDM-914A (4.610), CML-474 × V-351(-4.396), KDM-927 × V-351 (-3.110), KDM-930 × V 335(2.896) for wide variety of kernels rows per cob, KDM-347 × V-351(-7.544), KDM-440 × V-351(6.438), KDM-916A × KDM-914A (6.117), CML-470 × V-351 (5.992), CML-474 × V-335(5.274) for Number of kernels per row, KDM-340 × V-351 (3.966), KDM340 × KDM-914 A (3.150), KDM-930 × V-351(3.165), KDM-440 × KDM-914 A(3.556), KDM-347 × V-351(3.324) for 100 grain weight, KDM-347 × V351 (34.197), KDM-440 × V335(28.933), CML470 × KDM-914A (29.380), KDM-895 × KDM914A (25.383), KDM927A × KDM-914 (19.549) for Grain yield per plant. Significant heterosis for yield and yield related constituent qualities and most flowering

traits have been said by way of a variety of people [5,6], Alam et al., 2008; [7,8], Jawaharlal et al., 2012; Ali et al., 2012; [9,10] and Singh et al., 2013).

4. CONCLUSION

Out of 10 diverse lines maximum grain yield was recorded in CML470 × KDM-914 A ranging from 60 q/hawith maximum 100 grain weight of 22(g) followed by CML474 × V351 with grain yield of about 58 q/ha and least grain yield in KDM-347x V351 and KDM-340 × V351which was about 13 q/ha. So further research can be undertaken using these hybrids for crossing purpose CML470 × KDM-914 and CML474 × V351.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Dar SA, Ali G, Rather AG, Khan MN. Combining ability for yield and maturity traits in elite inbred lines of

- maize (*Zea mays* L.) International Journal of Agricultural Sciences. 2017;3(2):290-293.
2. Dar ZA, Lone AA, Alaie BA, Dar SA, Ali G, Abidi I, Lone RA, Gulzar S, Yousuf N. Genetic variation in maize inbred lines under temperate conditions. Journal of Experimental Biology and Agriculture Sciences. 2017;5(2):215-219
 3. Dar ZA, Lone AA, Alaie BA, Gowhar A, Asima G, Abidi I. Estimation of combining ability involving quality protein maize (QPM) inbreds under temperate conditions. The Bioscan. 2015;10(2): 863-867.
 4. Dar ZA, Lone AA, Khuroo NS, Ali G, Abidi I, Ahangar MA, Wani MA, Yasin AB, Gaza A, Lone RA, Yousuf N, Gulzar S. Line x tester analysis in maize (*Zea mays* L.) for various morpho-agronomic traits under temperate conditions. International Journal of Current Microbiology and Applied Sciences. 2017;6(7):1430-1437.
 5. Jain R, Bharadwaj DN. Heterosis and inbreeding depression for grain yield and yield contributing characters in quality protein maize. Agricultural Communications. 2014;2(1):8-16.
 6. Rajesh V, Kumar SS, Reddy VN, Sankar AS. Heterosis studies for grain yield and its component traits in single cross hybrids of maize (*Zea mays* L.). International Journal of Plant, Animal and Environmental Sciences. 2014;4(1):304-306.
 7. Amiruzzaman M, Islam MA, Hassan L. Rohman MM. Combining ability and heterosis for yield and component characters in maize. Academic Journal of Plant Sciences. 2010;3(2):79-84.
 8. Amanullah, Jehan S, Mansoor M, Khan MA. Heterosis studies in diallel crosses of maize. Sarhad Journal of Agriculture. 2011;27(2):207-211.
 9. Abuali AI, Abdelmulla AA, Khalafalla MM, Idris AE, Osman AM. Combining ability and heterosis for yield and yield components in maize (*Zea mays* L.). Australian Journal of Basic and Applied Sciences. 2012;6(10):36-41.
 10. Izhar T, Chakraborty M. Combining ability and heterosis for grain yield and its components in maize inbreds over environments (*Zea mays* L.). African Journal of Agricultural Research. 2013; 8(25):3276-3280.

© 2021 Viqar-Un-Nisa et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/79633>