



Productivity of Aerobic Rice under Different Lateral Arrangement and Nutrient Management

**Sanjay Kumar ^{a*}, Sanjay K. Dwivedi ^{a#}, Sarjeet Singh ^b, Gharsiram ^c
and Pradeep Kumar ^c**

^a Department of Agronomy, IGKV, Raipur (Chhattisgarh)-492012, India.

^b Department of Extension Education, SKNAU, Jobner (Rajasthan)-303329, India.

^c Department of Agronomy DRPCAUPusa Samastipur (Bihar)-848125, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2021/v33i2430797

Editor(s):

(1) Dr. Ahmed Medhat Mohamed Al-Naggar, Cairo University, Egypt.

Reviewers:

(1) B. S. Gohil, Junagadh Agricultural University, India.

(2) K. Vanitha, Tamil Nadu Agricultural University, India.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/79343>

Original Research Article

Received 09 October 2021

Accepted 19 December 2021

Published 21 December 2021

ABSTRACT

An experiment to assess the "Productivity and economic feasibility of lateral arrangement and nutrient management in aerobic rice" was conducted at the Instructional-cum-Research Farm, I.G.K.V, Raipur, (C.G.) during *kharif* season of 2018. The soil of experimental field was clayey (*Vertisols*) in texture. The trial was laid out in strip plot design *viz.*, horizontal strips consist of 3 lateral arrangements and vertical strips consist of 4 nutrient management with 12 treatment combinations. The treatments consisted of 3 lateral arrangements *viz.*, M₁- lateral at 25 cm (1 LPH/0.3 metre spacing), M₂- lateral at 50 cm (2 LPH/0.3 metre spacing) and M₃- conventional practice and 4 nutrient management of N₁- 50 % RDF, N₂ - 100 % RDF, N₃ - 150 % RDF and N₄ - STCR- based fertilizer recommendation. Aerobic rice variety Indira aerobic -1 was used for trial. The sowing was done on 20th June, 2018 at 20 cm of spacing and harvesting of crop was done on 7th October, 2018.

The results of trial revealed that the growth parameters like plant height (cm), number of leaves hill⁻¹, dry matter accumulation hill⁻¹ (g), number of tillers (m⁻²) and grain yield (t ha⁻¹) were significantly higher under lateral arrangement at 25 cm (M₁). However, lowest value of all these parameters

[#]Senior Scientist;

^{*}Corresponding author: E-mail: sanjaygugrwal1995@gmail.com;

were obtained in conventional practice (M₃). Similarly, all these characters were also higher among nutrient management with application of STCR based fertilizer (N₄). Remarkably, lowest values were obtained with application of 50 % RDF (N₁).

Keywords: *Aerobic rice; STCR; conventional; yield; test weight.*

1. INTRODUCTION

Rice (*Oryza sativa* L.) crop is the most important staple cereal crop of more than 65% population in Asia and it comes in the second position among the food crops after wheat. This grain crop provides 20 percent of world's dietary energy supply, while maize and wheat crops supplies 5 and 19 percent, respectively. In Asia, more than 2 billion people are consuming about 60-70 % of their energy requirement and derived products from rice crop [1]. It covers more than 85 per cent of total production that is consumed by human beings however, it deserves a prime place among cereals as the world's most important wetland crop.

With the scarcity of water resources and competition with other sectors, the contribution of water for irrigation purpose is decreased by 10 to 15 per cent in the next 2 decennium. Over the past decade, In globally we are suffering for water scarcity and competition for water. As the water demand for municipal, domestic, environmental and industrial purposes rises in the near future, thus water availability for agriculture sector gets affected. Water status for agriculture purposes is estimated at about 83.3 per cent of the total water used today, will shrink to 71.6 per cent in 2025 and to 64.6 per cent in 2050 [2]. Rice is a high water requirement crop which requires 3000- 5000 litres of water to produce one kg of grain which is almost 2 to 3 times higher than any other cereal crops like wheat, maize [3]. In this case, One of the ways that we need to increase water use efficiency. Several techniques of water saving have been developed for rice crop such as saturated soil culture, alternate wetting and drying method [4], system of rice intensification and raised bed system to lower the water requirements of the rice crop. Basically, Rice prefers more water and flooded condition, but recent technologies and developments demonstrate showing that rice can be also grown in dry soils under less water conditions which is called aerobic rice. The water productivity of rice under aerobic conditions was 32-88% higher than under flooded conditions. Aerobic rice is defined as a production system in which, direct seeding of high yielding and input

responsive rice cultivars with aerobic adaptation has grown in non-puddled, non-flooded and non-saturated soil during the entire growing cycle.

Drip irrigation, also known as trickle irrigation is an irrigation method that applies water slowly to the root zone of plants, through a network of valves, pipes, tubes and emitters. The goal is to optimize water and input usage. Application of water-soluble fertilizers (WSF) through irrigation as a carrier improving nutrient use efficiency. Thus nutrients in the irrigation water will likely be placed directly to the root zone of crop [5]. Fertigation is the precise application of water soluble fertilizers through sprinkler and drip irrigation. It is an efficient and agronomically sound method of providing soluble plant nutrients directly to the active plant root zone. Fertigation is a new agricultural technique, which supplies water and fertilizer simultaneously [6,7]. It can supply fertilizer and water at the right time and right place thus, improving water-use efficiency and uptake of nutrients.

Drip fertigation permits the application of nutrients directly at the site of high concentration of active roots. Since nutrients are applied to a limited soil volume, the fertilizer use efficiency is also high. Adoption of micro irrigation might help in increasing the irrigated area, productivity of crops, water use efficiency and also achieve more weed control efficiency (83 %) by making non-availability of irrigation water to the weeds [8]. On the other hand, conventional fertilization especially on light soils may cause huge nutrient losses through leaching, percolation and volatilization.

2. MATERIALS AND METHODS

The field experiment was conducted during *kharif season* from 20 June 2018 to 7th October 2018 at Research cum Instructional Farm, Indira Gandhi Krishi Vishwavidyalaya, Raipur, Chhattisgarh. The experimental site is situated at 21°4' N latitude and 81°35' E longitude at the height of 290.20 m above the mean sea level (MSL). According to the data, aerobic rice crop received 964.5 mm rainfall during the crop period from 20th June 2018 to 7th October 2018 and

maximum temperature was varied from 31.2°C to 37.24 °C, while minimum temperature varied from 24.53 °C to 27.13°C. Relative humidity throughout the crop season during 2018 was varied between 70 to 95 percent. The open pan evaporation mean values ranged from 21.50 to 28.0 mm day⁻¹ whereas, average sunshine hours varied from 1.8 to 8.36 hours day⁻¹. The average wind velocity varied from 0.88 to 8.01 km hr⁻¹ of crop season during 2018 for different weeks. Soil type of the experimental field is texturally clay, *Vertisols* and locally known as “*kanhar*”. Indira aerobic-1 variety of aerobic rice maturing in about 115-120 days was sown with a spacing of 20 cm in rows. The experiment was consisted of 12 treatments with three replications. The treatments were divided into horizontal and vertical strip with strip plot design. The horizontal strips were divided into two lateral arrangements and one conventional method of irrigation and vertical strip was further divided into four fertigation levels. Horizontal strips consisted, M₁- Lateral at 25 cm (1 LPH/0.3 Meter spacing), M₂- Lateral at 50 cm (2 LPH/0.3 Meter spacing), M₃- conventional practice (surface irrigation) and Vertical strips consists N₁- 50 % RDF, N₂- 100 % RDF (100:60:40) kg ha⁻¹ (N:P₂O₅:K₂O), N₃- 150 % RDF, N₄- STCR based fertilizer recommendation (for 7 tonnes). The sowing was taken up on June 20th, 2018 by maintaining the intra and inter row spacing of 20 cm with seed rate of 80 kg ha⁻¹. Immediately after sowing the seeds are covered with soil with the help of planking. Water soluble fertilizers used were Urea, Phosphoric acid (PA) and Murate of potash (MOP) as N, P and K sources, respectively. The experimental data recorded were subjected to statistical analysis by using Fisher's method of Analysis of Variance (ANOVA). The level of significance used in the F and t-test was a p=0.05.

3. RESULTS AND DISCUSSION

3.1 Growth Attributes

The growth attributes of aerobic rice are influenced by lateral arrangement and nutrient management and are presented in Table 3. Among lateral arrangements, lateral at 25 cm (1 LPH/0.3 meter distance, M₁) recorded higher plant height, number of tillers m⁻², total dry matter production hill⁻¹, number of leaves hill⁻¹, Leaf area hill⁻¹, Leaf area index, Plant population (114.58 cm, 495.83 m⁻² and 44.13 g hill⁻¹, 40.50, 1094.67, 5.47, 80.83 respectively). This is might be due to the application of water and nutrients through

drip fertigation directly to the root zone make them easily available and also maintaining moist conditions around root zone, which creates a favourable environment for roots to grow and absorb water and nutrients more effectively help in increasing number of tillers, leaves, total dry matter production and increasing plant height. These results are in conformity with the findings of Hebbar *et al.* [9] and Gururaj [10]. Among nutrient management significantly higher plant height, number of tillers m⁻², total dry matter production hill⁻¹ number of leaves hill⁻¹, Leaf area hill⁻¹ and Plant population were observed with the application of STCR based fertilizer recommendation (N₄) (113.56 cm, 519.33 m⁻², 45.58 g hill⁻¹, 42.89, 1108.48, 80.0 respectively). This is mainly because application of fertilizers through drip irrigation resulted in continuous supply of nutrients besides maintaining optimum water availability which leads to higher uptake of nutrients which in turn recorded higher growth attributes [10]. Further, significantly higher plant height, number of tillers m⁻², total dry matter production hill⁻¹, number of leaves hill⁻¹, Leaf area hill⁻¹, Leaf area index and Plant population in fertigation treatments over surface irrigation resulted in the production of higher plant biomass that could be attributed for higher plant growth parameters under drip fertigation treatments. Such finding is in conformity with the findings of Hebbar *et al.*, [9] in tomato; Vijaykumar [11] in rice and Soman, [12] in rice. However, lower values of these parameters observed under conventional practices (M₃) and application of 50 % RDF.

A significant difference in the number of tillers m⁻² (at harvest) was observed due to the interaction of lateral arrangement and nutrient management and data are presented in Table 2. Lateral at 25 cm (1 LPH/0.3 meter distance, M₁) with the application of 150 % RDF (N₃) has recorded significantly higher number of tillers, which was statistically at par with Lateral at 25 cm (1 LPH/0.3 meter distance, M₁) with the application of STCR based fertilizer (N₄) and Lateral at 50 cm (2 LPH/0.3 meter distance, M₂) with N₃ and N₄ treatment. This is might be due to application of water and nutrients directly to the root zone make them easily available and also maintaining moist conditions around root zone, which creates favourable environment for roots to grow and absorb water and nutrients more effectively help in increasing number of tillers and significantly lower number of tillers was observed under conventional practices (M₃) with the application of 50 % RDF (N₁).

Table 1. Interaction between lateral arrangement and nutrient management on Test weight (g) of aerobic rice

Lateral arrangement	Nutrient management				Mean
	N ₁	N ₂	N ₃	N ₄	
M ₁ : Lateral at 25 cm	20.56	23.28	23.62	23.85	22.83
M ₂ : Lateral at 50 cm	20.14	20.24	22.13	23.48	21.50
M ₃ : Conventional practice	19.68	20.60	21.69	22.06	21.01
Mean	20.13	21.37	22.48	23.13	
				SEm±	CD (P=0.05)
Two horizontal strip means at the same level of vertical strip				0.42	1.43
Two vertical strip means at the same level of horizontal strip				1.02	3.49

Table 2. Interaction between lateral arrangement and nutrient management on No. of tillers m⁻² (At harvest) of aerobic rice

Lateral arrangement	Nutrient management				Mean
	N ₁	N ₂	N ₃	N ₄	
M ₁ : Lateral at 25 cm	437.67	451.00	559.33	535.33	495.83
M ₂ : Lateral at 50 cm	412.67	443.00	526.67	535.33	479.42
M ₃ : Conventional practice	399.67	431.33	432.67	487.33	437.75
Mean	416.67	441.78	506.22	519.33	
				SEm±	CD (P=0.05)
Two horizontal strip means at the same level of vertical strip				18.63	65.8
Two vertical strip means at the same level of horizontal strip				14.39	46.0




Fig. 1. A general view experimental site of aerobic rice as influenced by lateral arrangement and nutrient management

Table 3. Growth attributes of aerobic rice influenced by lateral arrangement and nutrient management

Treatment	Plant height (cm) at harvest	No. of tillers (m ²) at harvest	Total dry matter production (hill ⁻¹) at harvest	No. of leaves hill ⁻¹ at harvest	Leaf area hill ⁻¹ (cm ²) at 90 DAS	Leaf area index at 90 DAS	Plant population (m ⁻²)
Lateral Arrangement							
M ₁ : Lateral at 25 cm (1LPH/0.3 MD)	114.58	495.83	44.13	40.50	1094.67	5.47	80.83
M ₂ : Lateral at 50 cm (2LPH/0.3 MD)	109.50	479.42	38.58	37.33	1032.11	5.16	75.58
M ₃ : Conventional practice	104.25	437.75	31.83	33.08	950.01	4.75	74.00
SEm ±	1.86	13.68	0.70	0.55	25.69	0.12	1.29
CD (P=0.05)	7.30	53.73	2.75	2.19	100.89	0.50	NS
Nutrient Management							
N ₁ : 50 % RDF	103.44	416.67	28.04	26.00	904.82	4.52	73.00
N ₂ : 100% RDF	107.22	441.78	37.18	37.67	975.15	4.88	75.22
N ₃ : 150 % RDF	113.56	506.22	41.91	41.33	1113.94	5.57	79.00
N ₄ : STCR based (for 7 tonnes)	113.56	519.33	45.58	42.89	1108.48	5.54	80.00
SEm ±	2.07	8.07	0.75	1.89	27.96	0.14	0.90
CD (P=0.05)	7.16	27.93	2.58	6.53	96.77	0.48	NS
Int (WXN)	S	S	S	S	S	S	NS

*MD = Distance of emitter (in meter)

Table 4. Yield and Yield Parameters of aerobic rice influenced by lateral arrangement and nutrient management

Treatment	Grain yield (t ha ⁻¹)	No. of panicles m ⁻²	No. of grains panicle ⁻¹	Test weight (g)	Sterility (%)
Lateral Arrangement					
M ₁ : Lateral at 25 cm (1LPH/0.3 MD)	6.35	314.58	177.58	22.83	14.02
M ₂ : Lateral at 50 cm (2LPH/0.3 MD)	5.96	265.83	163.25	21.50	15.28
M ₃ : Conventional practice	4.93	226.08	134.00	21.01	13.73
SEm ±	0.09	13.42	1.71	0.26	0.60
CD (P=0.05)	0.38	52.70	6.75	1.04	NS
Nutrient Management					
N ₁ : 50 % RDF	4.97	199.44	125.22	20.13	16.56
N ₂ : 100% RDF	5.68	231.22	139.56	21.37	15.10
N ₃ : 150 % RDF	6.14	300.89	173.89	22.48	12.49
N ₄ : STCR based (for 7 tonnes)	6.19	343.78	194.44	23.13	13.22
SEm ±	0.11	11.81	3.27	0.97	0.84
CD (P=0.05)	0.40	40.86	11.33	NS	2.90
Int (WXN)	S	S	S	S	S

*MD = Distance of emitter (in meter)

3.2 Yield and Yield Parameters

Yield contributing characters of aerobic rice viz., total no. of panicles m^{-2} , total number of grains panicle⁻¹, Test weight (1000 grain weight in gm) were found significantly higher except test weight influenced by nutrient management (Table 3). The significant result was found in grain yield of aerobic rice (Table 4). Among lateral arrangements, lateral at 25 cm (1 LPH/0.3 meter distance, M₁) recorded higher grain yield (6.35 t ha⁻¹). It is 28% higher than conventional practice and lower values of grain yield was found in conventional practices (M₃). Drip fertigation in aerobic rice improved the availability of nutrients in the root zone for plant uptake leading to better growth and development of plants and dry matter accumulation. The improved dry matter production enhanced grain yield in drip fertigation treatments. Under Closer lateral spacing better prevalence of soil moisture and nutrients are more easily available to plant than broader lateral spacing. Hence, better availability of nutrients under M₁ led to more uptake and better growth and development of plants which resulted in remarkably higher grain yield of aerobic rice.

Among nutrient management, Higher grain yield (6.19 t ha⁻¹) was found with the application of STCR based fertilizer (N₄) which is 24% higher than application of 50% RDF (N₁). Higher grain yield may be due to its superiority in producing higher productive tillers hill⁻¹, No. of panicles hill⁻¹, 1000-seed weight and total number of filled grains panicle⁻¹ with lower values of sterility percentage than the other treatment. The lower values of grain yield were registered with the application of 50 % RDF (N₁). The highest yield and yield attributes in STCR might be due to the fact that soil testing helps the farmers to use fertilizers according to needs of crop. Fertilizer use for targeted yield is an approach, which takes into account the crop needs and nutrients present in the soil. The treatment with STCR based fertilizer application realized the target yield and closely accorded with those reported by Ray *et al.*, (2000) ref missing, Meena *et al.*, [13] and Ramesh and Chandrashekar, (2007).

Significant difference on test weight (gm) was observed due to the interaction of lateral arrangement and nutrient management and data are presented in Table 1. Lateral at 25 cm (1 LPH/0.3 meter distance, M₁) with the application of STCR based fertilizer (N₄) was recorded significantly highest test weight, which was statistically at par with all combination of

treatment except Lateral at 50 cm (2 LPH/0.3 meter distance, M₂) with (N₁), (N₂) and conventional practices (M₃) with (N₁) treatments. The lower test weight was observed under conventional practices (M₃) with the application of 50 % RDF (N₁).

4. CONCLUSION

Drip fertigation permits the application of nutrients directly at the site of high concentration of active roots. Since nutrients are applied to a limited soil volume, the fertilizer use efficiency is also high. Conventional fertilization especially on light soils may cause huge nutrient losses through leaching, percolation and volatilization.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Geethalakshmi V, Ramesh T, Azhagu P, Lakshmanan A. Agronomic evaluation of rice cultivation systems for water and grain productivity. *Archi. Agron. Soil Sci.*, 2011;57(2):159-166.
2. Yadav JSP. Conservation and managing water resource for sustainable agriculture. *J. Water Manage.* 2002;10(1-2):1-10.
3. Cantrell RP, Hettel GP. New challenges and technological opportunities for rice based production systems for food security and poverty alleviation in Asia. Presented at the Rice Conf., FAO, Rome, Italy held on 12-13 February. 2004;2004:247-251.
4. Tabbal DF, Bouman BA, Safdos MA. On farm strategies for reducing water input in irrigated rice. *Agric. Water Mng.* 2002;56(2):93-112.
5. Clark GA, Stanley CD, Maynard DN, Hochmuth GJ, Hanlon EA, Haman DZ. Water and fertilizer management of micro irrigated fresh market tomatoes. *American Soc. Agri. Engineering.* 1991;34: 429-435.
6. Castellanos MT, Tarquis AM, Ribas F, Cabello MJ, Arce A, Cartagena MC. Nitrogen fertigation: an integrated agronomic and environmental study. *Agricultural Water Management*;2012.
7. Mahajan G, Chauhan BS, Timsina J, Singh PP, Singh K. Crop performance and water and nitrogen use efficiencies in dry-seeded rice in response to irrigation and fertilizer amounts in Northwest India. *Field Crops Research.* 2006;134:59-70.

8. Sivanappan RK. Irrigation and rain water management for improving water use efficiency and production in cotton crop. In: Proceedings of International Symposium on "Strategies for Sustainable Cotton Production a Global Vision. Univ. Agric. Sci., Dharwad, Karnataka;2004.
9. Hebbar SS, Ramachandrappa BK, Nanjappa HV, Prabhakar M. Studies on NPK drip fertigation in field grown tomato (*Lycopersicon esculentum* Mill.). European J. Agron. 2004;21:117-127.
10. Gururaj K. Optimization of water and nutrient requirement through drip fertigation in Aerobic rice;2013.
11. Vijaykumar P. Optimization of water and nutrient requirement for yield maximization in hybrid rice under drip fertigation system. M.Sc. (Agri.) Thesis (Unpub.), Tamil Nadu Agric. Univ., Coimbatore;2009.
12. Soman P. Drip fertigation for rice cultivation, In: Proc. Asian Irrig. Forum, 11-12 April, 2012, ADB, Manila, Philippines; 2012.
13. Meena M, Ahmed S, Riazuddin M, Chandrasekhara KR, Rao BRCP. Soil test crop response calibration studies on onion (*Allium cepa*) in alfisols. Journal of Indian Society of Soil Science. 2001;49:709–713.

© 2021 Kumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/79343>