



Effect of Foliar Application of Gibberellic Acid on Plant Growth, Flowering and Yield Attributes in Pansy (*Viola × wittrockiana* Gams.)

**Anu Sharma ^{a*}, B. S. Dilta ^b, C. L. Sharma ^c, Manish Kumar ^d,
Akhilesh Koundal ^a and Shivani Sharma ^d**

^a Department of Seed Science and Technology, Dr. Y S Parmar University of Horticulture and Forestry, Nauni, Solan-173230, India.

^b Department of Floriculture and Landscape Architecture, Dr. Y S Parmar University of Horticulture and Forestry, Nauni, Solan-173230, India.

^c Department of Fruit Science, Dr. Y S Parmar University of Horticulture and Forestry, Nauni, Solan-173230, India.

^d Department of Vegetable Science, Dr. Y S Parmar University of Horticulture and Forestry, Nauni, Solan-173230, India.

Authors' contributions

This work was carried out in collaboration among all authors. Authors AS and BSD designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript, managed the analyses of the study. Authors CLS and MK managed the literature searches and guides during the research. Authors AK and SS helps in laboratory work. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i173222

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/101971>

Original Research Article

Received: 18/04/2023

Accepted: 20/06/2023

Published: 09/07/2023

*Corresponding author: E-mail: anusharma5193@gmail.com;

ABSTRACT

An investigation was carried out to study the effect of foliar application GA₃ on plant growth and quality attributes in Pansy (*Viola × wittrockiana* Gams.) during 2019-2020. The field experiment was laid out at randomized complete block design (RCBD). The 18 treatment combinations including 6 doses of GA₃ *i.e.*, 50, 100, 150, 200, 250 and 300 ppm were applied at 3 durations *i.e.*, 30, 40 and 50 DAT along with a non-factorial control. Of them, GA₃ @ 300 ppm at 40 DAT showed significant increment in plant height (34%), plant spread (32%), number of flowers per plant (25%), number of flowers per plot (25%), size of flowers (35%), flowering duration (25%), seed yield per plant (68%), seed yield per plot (68%) and seed yield per hectare (68%), number of seeds per capsule (48%), number of capsules per plant (38%) as compared to control checks whereas earliest flowering was observed due to GA₃ @ 300 ppm when applied at 30 DAT.

Keywords: Foliar; GA₃; pansy; RCBD; DAT.

1. INTRODUCTION

Pansy (*Viola × wittrockiana* Gams.), a popular, commercially important ornamental bedding plant in both developed and developing countries [1]. It is a winter annual and belongs to family Violaceae. The word 'Pansy' has been derived from a French word 'Pensee' meaning 'loving thoughts' [2]. It is basically derived from *Viola tricolor*. Pansy is native to Central Europe with abundant flower colors [3,4]. The Pansy producing purple colour flowers is believed to represent sweet memories. It is well-documented that the origin of modern pansy (*Viola × wittrockiana* Gams.) began in England in 1815 by hybridizing wild pansy (*Viola tricolour* L.) with *Viola lutea* and *Viola altaica* [5]. Pansies are divided in to two groups *viz.*, 'Clear types' bearing flowers with solid colour and 'Faced type', which are multi-coloured or with spots or patches of various colours. However, based on flower size, the pansy is categorized in three main types namely, large (with a bloom of 7.5-10 cm diameter), medium (bloom size 5.0-7.5 cm), and small (bloom size of 2.5-5.0 cm), respectively.

The pansy flowers are borne singly on the long stalk with attractive colours and shapes. These are ideal for utilizing in winter landscapes as they are cold tolerant plants and grow more luxuriantly with profuse flowering under cool and moist environment. At present, pansy is one of the dominating plant species among ornamental bedding plants in the U.S. market and, various breeding programs has been performed in Germany, U.S. and Japan to develop novel colors and for other characteristics [6]. They are often cultivated in garden beds, pots, borders, hanging baskets, or landscapes as annuals or biennials [7].

The plant growth regulators of either natural or synthetic origin may incite some changes in the various processes of growth, development and flowering of plants in general and ornamentals in particular by manipulating the requisite physiological, biochemical and flower induction reactions during primary and secondary metabolism [8]. Gibberellins are naturally occurring plant growth regulators that are produced in response to both developmental and environmental cues [9]. Yuan *et al.*, [10] has reviewed effect of GA₃ and stated that it regulates plant height, more number of branches as well as increase productivity. Gibberellins have an impact on plants at every phase of their life cycles, showing a remarkable diversity in their physiological activities [11].

Numerous research works have revealed the effects of GA₃ on plant growth, flowering and yield attributes in several crops. Shrestha *et al* [12], have shown that foliar application of gibberellic acid significantly improves quality of calendula flowers. Biswas *et al.*, [13], observed better plant growth and yield attributing characters with application of GA₃. Khangjarakpam *et al* [14], also observed similar results in marigold with application of GA₃ in China aster. Husain *et al* [15] also observed increase in seed yield with GA₃ application in pea.

2. MATERIALS AND METHODS

The experiment was conducted on pansy (*Viola × wittrockiana* Gams.) during winter season of 2019-2020 at Dr YS Parmar University, Nauni, Solan. The treatment combinations consisted of 6 different doses of gibberellic acid (GA₃), *viz.*, 50 ppm, 100 ppm, 150 ppm, 200 ppm, 250 ppm and 300 ppm at three different times *i.e.*, 30

DAT, 40 DAT and 50 DAT along with control. The experimental design followed in field was randomized complete block design (RCBD) with three replications per treatment. The plant spacing was 30 × 25 cm which occupies 12 plants per plot. Sowing was done on first fortnight of October and transplanting was carried out 30 days after sowing and standard cultural practices like hoeing, irrigation, weeding and fertilizer application were followed as per the requirement of the crop. Data on various growth, flowering and yield parameters were recorded and statistically analyzed by applying the technique of analysis of variance. Statistical interference was implied with ANOVA, keeping the level of significance at 5% (P=0.05).

3. RESULTS AND DISCUSSION

3.1 Growth Parameters

The analysis of variances obtained for the parameters under study have exhibited significant variations among different treatments alone and in combinations as well. As regards the effects of GA₃ doses, G₆ (GA₃ @ 300ppm), recorded highest values for plant height (30.24 cm). Moreover, GA₃ application at 40 DAT (40 days after transplanting; D₂) produced maximum plant height (27.44 cm). Maximum plant height (31.29 cm) was significantly induced by the interaction effects of G₆ × D₂. In general, the increase of plant height by GA₃ treatment could be due to the induction of cell division and cell elongation on faster rate. The results are in line with the research findings obtained in African marigold cultivar's 'Pusa Narangi' and 'Pusa Basanti' [16], *Dahlia pinnata* [17], sunflower [18] in as well as *Calendula officinalis* cv. 'Bon Bon' [19].

The plant spread was also found higher in G₆ (25.81 cm), D₂ (24.11 cm) and their interaction G₆ × D₂ (17.87), which could be as a consequence of division and enlargement of cells and tissues as well as production of more number of side branches. Similar results have been reported by Khudus et al [19] in *Calendula officinalis* cv. 'Bon Bon' and Khangjarakpam et al [14] in African marigold.

More number of flowers per plant, flowers per plot, is higher when GA₃ is applied @ 300 ppm as well as when applied at 40 days after transplanting (See Tables 1 & 2). Their interaction G₆ × D₂ produced highest flowers (See Table 3). The production of more flowers

per plant might be as a consequence of producing maximum number of lateral particularly at juvenile stages of growth leading to developing of maximum vegetative buds. In due course of time, these buds have differentiated and converted in to flowering buds. Similar results have also been documented in gerbera [20], Sweet William [21], and marigold cv. 'Pusa Narangi Gaiinda' [22].

Increase in size of flowers (2.70), is when 300 ppm GA₃ is applied as well as when applied at D₂ i.e., 40 DAT (2.24 cm). The interaction G₆ × D₂ significantly shows increase in the size of flower (2.81 cm). The application of GA₃ particularly at higher doses and optimum stage of growth might have increased flower size as a consequence of forming a better source - sink duo so as to accumulate higher concentration of metabolites and their utilization by the plants resulting in the production of large size flowers. Similar results have also been documented in chrysanthemum [23], carnation [24], *Dianthus caryophyllus* L [25], calendula [12].

The maximum duration of flowering (88.00 days) occurs with application of 300 ppm of GA₃. The increase in flowering duration results in D₂ (83.50 days). The interaction G₆ × D₂ significantly shows extend in flowering duration (89.33 days).

3.2 Seed Yield and Yield Contributing Characters

More number of capsules per plant (53.33) were observed to be significantly superior in G₆. The number of capsules per plant (47.39) is higher in D₂ (GA₃ at 40 DAT). The interaction G₆ × D₂ significantly produced more number of capsules (54.33) per plant. More number of capsules per plant could be attributed to the production of maximum flowers of better size and quality. Subsequently, these flowers upon pollination could develop capsules in sufficient quantum, hence, more number of capsules were produced per plant.

More number of seeds per capsule (64.00) were significantly higher in GA₃ i.e. 300 ppm (G₆). The number of seeds per capsule (52.50) was higher with application of GA₃ at 40 DAT. The interaction G₆ × D₂ significantly produced more number of seeds (52.50) per capsule. Production of more seeds with higher doses of GA₃ at proper stage may be as a consequence of better growth and flowering of plants as well as improving pollination and fertilization resulting in better

Table 1. Effect of different doses and durations of GA₃ on various growth and quality parameters of pansy (*Viola × wittrockiana* Gams.)

Treatments	Plant height (cm)	Plant spread (cm)	Flower diameter (cm)	Days to first flowering	Duration of flowering	Number of flowers /plant	Number of flowers /plot	Number of capsules/ plant	Number of seeds/capsule	Seed yield /plant (g)	Seed Yield/ plot (g)	Seed Yield/ ha (kg)
Doses of GA₃ (ppm)												
G ₁ (GA ₃ 50ppm)	22.48	20.41	1.87	74.56	70.56	102.33	1228.00	38.67	37.11	1.44	17.25	172.47
G ₂ (GA ₃ 100ppm)	24.58	22.61	1.90	71.44	75.89	115.61	1387.33	41.67	44.78	1.87	22.18	221.83
G ₃ (GA ₃ 150ppm)	26.19	23.39	1.96	66.89	82.89	121.89	1462.67	44.89	47.33	2.13	25.50	255.01
G ₄ (GA ₃ 200ppm)	27.71	24.07	2.15	65.67	85.22	125.67	1508.00	47.89	52.44	2.51	30.15	301.53
G ₅ (GA ₃ 250ppm)	28.90	24.99	2.51	61.11	86.11	129.44	1542.22	52.11	60.33	3.14	37.74	377.39
G ₆ (GA ₃ 300ppm)	30.24	25.81	2.70	58.78	88.00	130.22	1562.67	53.33	64.00	3.42	41.00	409.95

Table 2. Effect of different durations of GA₃ application on various growth and quality parameters of pansy (*Viola × wittrockiana* Gams.)

Treatments	Plant height (cm)	Plant spread (cm)	Flower diameter (cm)	Days to first flowering	Duration of Flowering	Number of flowers /plant	Number of flowers /plot	Number of capsules/ plant	Number of seeds /capsule	Seed yield /plant (g)	Seed yield/ plot (g)	Seed yield /ha (kg)
Durations of GA₃												
D ₁ (30 DAT)	26.52	23.49	2.18	64.83	81.44	120.39	1444.67	46.33	51.33	2.43	29.14	291.36
D ₂ (40 DAT)	27.44	24.11	2.24	66.39	83.50	123.08	1477.00	47.39	52.50	2.54	30.35	303.53
D ₃ (50 DAT)	26.09	23.04	2.12	68.00	79.39	119.11	1423.78	45.56	49.17	2.29	27.42	274.20

Table 3. Effect of different doses and durations of GA₃ application on various growth and quality parameters of pansy (*Viola × wittrockiana* Gams.)

Treatments	Plant height (cm)	Plant spread (cm)	Flower diameter (cm)	Days to first flowering	Duration of Flowering	Number of flowers /plant	Number of flowers /plot	Number of capsules/ plant	Number of seeds/capsule	Seed yield /plant (g)	Seed Yield/ plot (g)	Seed Yield/ ha (kg)
T ₁ (Control)	20.63	17.87	1.82	79.67	67.33	98.00	1176.00	33.67	34.00	1.14	13.68	136.84
T ₂ G ₁ × D ₁	22.03	20.07	1.87	73.67	70.00	101.00	1212.00	38.67	37.33	1.44	17.34	173.36
T ₃ G ₁ × D ₂	23.63	21.20	1.89	74.00	72.00	105.67	1268.00	39.67	38.00	1.51	18.11	181.12
T ₄ G ₁ × D ₃	21.77	19.97	1.86	76.00	69.67	100.33	1204.00	37.67	36.00	1.36	16.29	162.92
T ₅ G ₂ × D ₁	24.33	22.93	1.90	70.00	76.33	115.00	1380.00	41.33	45.33	1.88	25.50	225.04
T ₆ G ₂ × D ₂	25.33	23.48	1.91	71.00	79.67	119.83	1438.00	42.67	46.33	1.98	23.05	230.53
T ₇ G ₂ × D ₃	24.07	21.42	1.89	73.33	71.67	112.00	1344.00	41.00	42.67	1.75	20.99	209.92
T ₈ G ₃ × D ₁	26.10	23.10	1.96	65.00	84.00	121.67	1460.00	44.67	47.33	2.12	25.38	253.84
T ₉ G ₃ × D ₂	26.87	24.13	1.98	67.67	85.33	123.67	1484.00	45.67	48.33	2.21	26.49	264.88
T ₁₀ G ₃ × D ₃	25.60	22.95	1.94	68.00	79.33	120.33	1444.00	44.33	46.33	2.05	24.63	246.32
T ₁₁ G ₄ × D ₁	27.67	24.10	2.17	64.00	85.00	125.67	1508.00	47.67	53.00	2.53	30.33	303.28
T ₁₂ G ₄ × D ₂	28.40	24.47	2.30	66.00	87.00	127.00	1524.00	48.67	54.33	2.65	31.74	317.40
T ₁₃ G ₄ × D ₃	27.07	23.63	1.99	67.00	83.67	124.33	1492.00	47.33	50.00	2.37	28.39	283.92
T ₁₄ G ₅ × D ₁	28.97	25.00	2.54	59.00	85.67	129.00	1548.00	52.67	60.33	3.18	38.13	381.32
T ₁₅ G ₅ × D ₂	29.13	25.23	2.56	61.33	87.67	131.00	1572.00	53.33	62.33	3.32	39.86	398.64
T ₁₆ G ₅ × D ₃	28.60	24.73	2.43	63.00	85.00	128.33	1506.00	50.33	58.33	2.94	35.22	352.20
T ₁₇ G ₆ × D ₁	30.00	25.75	2.65	57.33	87.67	130.00	1560.00	53.00	64.67	3.43	41.13	411.33
T ₁₈ G ₆ × D ₂	31.29	26.17	2.81	58.33	89.33	131.33	1576.00	54.33	65.67	3.57	42.86	428.60
T ₁₉ G ₆ × D ₃	29.43	25.52	2.63	60.67	87.00	129.33	1552.00	52.67	61.67	3.25	38.99	389.92

seed setting. Therefore, said treatment combination could produce maximum seeds in a capsule. These results are in line with the findings of Hoque and Haque [26], who observed more number of seeds per pod in mung bean with the application of higher doses of GA₃. Similarly Thakur *et al.* [22] in African marigold also observed more seeds per head.

The maximum seed yield (3.42 g/plant), (41.00 g/plot and (409.95 kg/ha) was found in G₆. The seed yield (2.29 g/plant), (27.42 g/plot) and (274.20 kg/ha) is highest in D₂. The interaction revealed that GA₃ @ 300 ppm after 40 days of transplanting G₆ × D₂ produced maximum seed yield (3.57 g/plant), (42.86 g/plot) and (428.60 kg/ha).

Whereas, the minimum plant height, plant spread, number of flowers per plant, number of flowers per plot, size of flowers, duration of flowering, seed yield, number of seeds per capsule, number of capsules per plant, as well as maximum time to first flowering were reported with the application of GA₃ @ 50 ppm (G₁; See Table 1).

However, the plant height, plant spread, number of flowers per plant, number of flowers per plot, size of flowers, duration of flowering, number of seeds per capsule, number of capsules per plant, seed yield were observed to be the minimum with the application of GA₃ after 50 days of transplanting (D₃; See Table 2).

On the contrary, the values for plant height, plant spread, number of flowers per plant, number of flowers per plot, size of flowers, duration of flowering, number of seeds per capsule, number of capsules per plant, seed yield, were observed to be the lowest in control (T₁; See Table 3).

4. CONCLUSION

Application of GA₃ had a beneficial influence on almost all growth and flowering characteristics of Pansy. Among all the six doses of GA₃, application of GA₃ @ 300 ppm was significantly superior over other five concentrations w.r.t. various growth, flowering, seed yield and seed quality parameters this could be due to the induction of cell division and cell elongation on faster rate which ultimately gives more lateral branches, higher flowers per plant and hence greater seed yield. The foliar spray of GA₃ at 40 DAT has exhibited its eminence over other durations in terms of various growth, flowering,

seed yield and seed quality parameters. Therefore it is concluded that GA₃ @ 300 ppm sprayed after 40 days of transplanting has proved to be the most effective treatment combination enhancing growth and yield parameters in pansy.

ACKNOWLEDGEMENT

I feel great contentment to express my profound regards and deep sense of gratitude to the esteemed teacher and advisor, Dr. BS Ditta, Principal Scientist, Department of Seed Science and Technology for his competent guidance, sustained encouragement, keen interest, meticulous suggestions, constructive criticism, perpetual assistance and painstaking efforts with his scientific knack during the entire course of investigations.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Gandolfo E, Hakim G, Geraci J, Feuring V, Giardina E, Di Benedetto A. Responses of Pansy (*Viola wittrockiana* Gams.) to the quality of the growing media. American Journal of Experimental Agriculture. 2016;12(3):1-10.
2. Desai BB. Seeds handbook: Biology, production, processing and storage. 2nd revised edition. New York: Marcel Dekker. 2009:376-377.
3. Chattopadhyay PK. Commercial Floriculture. Gene-Tech Books. New Delhi. 2007:236-257.
4. Wang T, Li J, Li T, Zhao Y, Zhou Y, Shi Y, Peng T, Song X, Zhu Z, Wang J. Comparative transcriptome analysis of anthocyanin biosynthesis in pansy (*Viola × wittrockiana* Gams.). Agronomy. 2022;12:919.
5. Hazar D, Boyar I, Dincer C, Ertekin C. Investigation of color and bioactive compounds of different colors from pansy (*Viola × wittrockiana* Gams.) dried in hot air dryer. Horticulturae. 2023;9:186p.
6. Du XH, Wang MG, Słomka A, Liu HC. Karyologic and heterosis studies of the artificial inter and intraspecific hybrids of *Viola × wittrockiana* and *Viola cornuta*. Hort Science. 2018;53:1300–1305.

7. Sharma R, Dilta BS, Baweja HS, Kumari P, Thaneshwari T, Sahare HA. Effect of seed priming and planting dates on growth, flowering and seed production of pansy (*Viola x wittrockiana* Gams.) Annals of Biology. 2018;34:176–180.
8. Heldet HW. Plant Biochemistry and Molecular Biology. Oxford University. Pres, London; 1997.
9. Iqbal M, Ashraf M. Gibberellic acid mediated induction of salt tolerance in wheat plants: growth, ionic partitioning, photosynthesis, yield and hormonal homeostasis. Environmental and Experimental Botany. 2013;86:76–85.
10. Yuan L, Wu X, Liao F, Ma G and Xu Q. Hybrid Rice Technology. China Agricultural Press, Beijing China. 2003:132.
11. Pradeep Kumar CM, Chandrashekar SY, Kavana GB and Supriya BV. A review on role and use of gibberellic acid (GA₃) in flower production. International Journal of Chemical Studies. 2020;8(1):3076-3084.
12. Shrestha B, Karki A, Shrestha J. Effect of foliar application of gibberellic acid (GA₃) on quality attributes of calendula flowers (*Calendula officinalis*) cv. Gitana Fiesta in Chitwan, Nepal. Journal of Agricultural Science and Practice. 2020;5(4):168-173.
13. Biswas S, Bordolui SK, Sadhukhan R. Response of China aster (*Callistephus chinensis* L.) genotypes towards foliar application of GA₃. American International Journal of Agricultural Studies. 2021;5(1):1-15.
14. Khangjarakpam G, Singh LJ, Maitra S and Mandal S. Influence of foliar application of gibberellic acid on growth, development, yield and biochemical constituents of African marigold cv. 'Pusa Narangi Gainda'. Journal of Pharmacognosy and Phytochemistry. 2019;8:1581-1585.
15. Husain AJ, Muhmood AG and Alwan AH. Interactive effect of GA₃ and Proline on nutrients status and growth parameters of pea (*Pisum sativum* L.). Indian Journal of Ecology. 2018;45(1):201-204.
16. Ramdevputra MV, Deshmukh HN, Bhutani AM, Savaliya JJ, Pansuriya AG and Kanzaria DR. Effect of different gibberellic acid (GA₃) concentrations on growth, flowering and yield of African marigold. The Asian Journal of Horticulture. 2009;4: 82-85.
17. Youssef ASM, Gomaa AO. Influence of GA₃ application and Kristal on fertilizer on growth, flowering and chemical composition of *Dahlia pinnata* plant (summer flowering type). In: 1st International Scientific Conference on Ornamentals Alexandria, held at Horticulture department, Faculty Agriculture, Moshtohor, Benha University, Egypt. 2008;1:191-207.
18. Sathy H, Patra SK, Mohanty CR. Effect of plant growth regulators on growth and flowering of ornamental sunflower. International Journal of Agricultural Science and Research. 2016;6:561-568.
19. Khudus S, Prasad VM, Jogdand SM. Effect of plant growth regulators on growth and flower yield of calendula cv. 'Bon Bon'. Chemical Science Review and Letters. 2017;6:1290-1294.
20. Patra SK, Beura S and Shasani T. Efficacy of GA₃ on growth and flowering regulation of *in vitro* raised hybrid gerbera under shade net. Agricultural Science Digest. 2015;35:173-177.
21. Thakur A, Dilta BS, Mehak DK, Sharma BP, Gupta RK. Influence of plant spacing and GA₃ on growth and flowering of sweet William (*Dianthus barbatus* L.). International Journal of Farm Sciences. 2018;8:122-125.
22. Thakur P, Dilta BS, Gupta YC, Mehak DK, Kumar P. Effect of planting dates, mulching and application of GA₃ on growth and flower yield of Marigold (*Tagetes erecta* L.) cv. 'Pusa Narangi Gainda'. International Journal of Current Microbiology and Applied Sciences. 2019;8:3028-3039.
23. Sharifuzzaman SM, Ara KA, Rahman MH, Kabir K, Talukdar MB. Effect of GA₃, CCC and MH on vegetative growth, flower yield and quality of Chrysanthemum. International Journal of Experimental Agriculture. 2011;2(1): 17-20.
24. Kumar V, Umara V, Singh M. Effect of GA₃ and IAA on growth and flowering of carnation. Hort Flora Research Spectrum. 2012;1(1):69-72.
25. El-Naggar AH, El-Naggar AAM, Ismaiel NM. Effect of phosphorus application and gibberellic acid (GA₃) on the growth and

- flower quality of *Dianthus caryophyllus* L. 26. Hoque M, Haque S. Effect of gibberellic acid on physiological contributing characters of mungbean. Journal of American-Eurasian Journal of Agricultural and Environmental Science. 2009;6(4): 400-410. Biological Sciences. 2002;5:401-403.

© 2023 Sharma et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/101971>