



Growth Performance of *Crescentia cujete* (Robx) Seedlings as Influenced by Different Watering Regimes

A. F. Aderounmu¹ and O. A. Ogidan^{1*}

¹*Federal College of Forestry, Forestry Research Institute of Nigeria, Ibadan, Nigeria.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CJAST/2019/v37i130273

Editor(s):

(1) Professor, Jerzy Nowacki, PhD, DsC West Pomeranian University of Technology, Szczecin Institute of Materials Science and Engineering Al. Piastow, Szczecin, Poland.

Reviewers:

(1) Junaid Iqbal, University of Agriculture, Faisalabad, Pakistan.

(2) Amina Rabe Musa, Usmanu Danfodiyo University, Nigeria.

(3) Dale Loussaert, USA.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50674>

Original Research Article

Received 03 June 2019

Accepted 15 August 2019

Published 21 August 2019

ABSTRACT

This study investigated the effects of different watering regimes on the growth of *Crescentia cujete*. The seedlings were subjected to six different watering regimes which include: W_1 – control (watering everyday), W_2 – watering once every two days, W_3 – watering once every three days, W_4 – watering once every four days, W_5 – watering once every five days and W_6 – watering once every six days; each treatment replicated six times. The experiment was arranged in a Completely Randomized Design (CRD). The study was carried out at the Central Nursery of Forestry Research Institute of Nigeria, Ibadan. Growth parameters assessed include: plant height (cm), collar diameter (mm), leaf production and the biomass accumulation which was sectioned into leaf, stem and the root. The data collected were subjected to one-way Analysis of Variance (ANOVA) and means separation was done using Duncan Multiple Range Test (DMRT) at 5% level of probability. The results showed that there were significant differences among the treatments in plant height and number of leaves produced while there was no significant difference in the collar diameter. However, seedlings watered once every 5 days (W_5) performed best in height (20.48 cm) and leaf production (18.42). It was revealed that there were significant differences in the leaf and stem dry

*Corresponding author: E-mail: ogidantosh@gmail.com;

weights among the treatments with W_5 having the highest weight while there were no significant differences in the root dry weight. *Crescentia cujete* optimized water shortage for its growth.

Keywords: *Crescentia cujete*; watering regime; growth; dry weight.

1. INTRODUCTION

Crescentia cujete commonly referred to as Calabash tree is a forest tree that belongs to the family of Bignoniaceae which originated from tropical America and later introduced to Africa [1]. According to Gilman and Watson [2], it grows to about 6 to 9 m in height with a wide crown and long branches covered with clusters of tripinnate leaves and gourd-like fruit. The large fruit has a thin hard shell and whitish pulp with dark-brown seeds that are thin and flat. The spoon-shaped leaves are 5 to 18 cm long and 2 to 5 cm broad and they are arranged in clusters along the stout twigs. It is frequently cultivated as ornamental tree.

Nearly all parts of the tree are utilized for diverse articles and equipment, the wood is used as tools handles, ribs in boat building and gourds for containers and musical instruments [3]. They are valued in the manufacture of handicrafts and musical instruments. The wood has specific gravity of 0.6 to 0.8; it is strong, flexible, moderately hard and heavy. Hence, it can be used for firewood and construction in rural areas and in the manufacture of handles for agricultural implements. The fruit pulp is laxative, emollient, expectorant and fever medicine. Fresh seeds are ground and mixed with water to make a refreshing drink which has sweet and pleasant taste [4].

Phenols and tannins are present in the fruit sample of *C. cujete*. Phenol and phenolic compounds have been extensively used in disinfections and remain the standard with which other bactericides are compared in official test [5,3]. This underscores why *C. cujete* is used as disinfectants and bactericides in emollient healing and in the treatment of burns [1]. Flavonoids found in *C. cujete* can act as anti-oxidants and protects the cells of the body from radical damage; free radicals are believed to damage cell and inflict various health-related problems [6].

Water, which is in a continual state of flux, is a very essential factor in the growth and development of plants [7]. Various vital

processes in plants such as cell division, cell elongation, stem as well as leaf enlargement and chlorophyll formation depends on plant water availability [8,9]. There is usually structural deformity in plants frequently leading to death when there is insufficient water below critical level [10]. The reduction in relative water contents affects physiological processes and hence plant growth (Awodola, 1984 unpublished MSc dissertation); [11].

Water is required by plants for the manufacture of carbohydrates and as a means for transportation of essential nutrients and minerals and cooling of plant leaves. About ninety-five percent of water in plants is used up in cooling via evapotranspiration while the remainder is utilised in physiological processes such as photosynthesis and respiration [12] and evapotranspiration rates increases with higher temperature [13,14]. However, water availability is seasonal coupled with climate change effects which manifests in longer dry spell or inundation giving way to waterlogging.

The water requirements of tree species differ due to their varying morphological and/or genetic makeup; this explains the disparate adaptive tendencies of different plants to water availability extremes such as water deficit and waterlogging [15]. Season-dependent water availability in nurseries spawns the need to study the responses of different tree seedlings in the nursery to different watering regimes in order to achieve optimum growth and engender effective water management.

The information about the response of *C. cujete* to different watering regime is sparse despite its diverse medicinal and domestic potentials. Hence, the objective of this study is to investigate the effects of different watering regimes on the early growth of *C. cujete* with a view to contributing to its conservation and efficient water management in the nursery.

2. MATERIALS AND METHODS

The experiment was carried out at the central nursery of Forestry Research Institute of Nigeria, Ibadan located on latitude 7°23' N and longitude

3°58'E. The climate is characterized by wet (April to October) and dry seasons (November to March). The annual rainfall ranging between 1300 - 1500 mm and annual mean relative humidity of 80-85% (Forestry Research Institute of Nigeria Annual Meteorological Report, 2015).

The experiment was carried out between January and March, 2017. The seeds of *C. cujete* were obtained from a mother tree at the herbal garden of Forestry Research Institute of Nigeria, Ibadan, Oyo State, Nigeria. The top soil (0-20 cm) was collected from the floor of a *Gmelina arborea* plantation. The seedlings were raised in a germination tray for two weeks, with sterilized river sand used as the growing media. The two weeks old seedlings were transplanted into polythene pots filled with 2 kg of top soil and watered to pot capacity with respect to the treatments. The growth parameters such as plant height (cm), collar diameter (mm) and leaf production were assessed weekly while the biomass estimation was carried out at the end of the growth assessment.

The height of the seedlings was measured with the aid of a measuring tape graduated in centimeters (cm) from the soil surface while the collar diameter was determined with the aid of a Vernier caliper.

Biomass estimation was carried out by selecting three (3) seedlings from each treatment. The seedlings were sectioned into leaves, stem and root and oven-dried at 70°C for 24 hours. The dry weight of the leaves, stem and root of the seedlings were weighed and recorded. The mean of the dry weights of the three seedlings selected in each treatment were calculated [16].

There were six treatments and each treatment was replicated six times making a total of thirty-six experimental units in all.

The treatments are:

- W₁ – control (watering everyday)
- W₂ – watering once every two days
- W₃ – watering once every three days
- W₄ – watering once every four days
- W₅ – watering once every five days
- W₆ – watering once every six days

The experimental design used was Completely Randomized Design (CRD). The data collected were subjected to a one-way Analysis of Variance (ANOVA) at 5% level of probability

while the means found to be significant were separated using Duncan Multiple Range Test (DMRT) at 5% level of probability [17].

3. RESULTS AND DISCUSSION

According to Federal Fertilizer Department of the Federal Ministry of Agriculture and Rural Development (2011), the soil was moderately acidic; the nitrogen and organic carbon contents were low while there was moderate amount of phosphorus and potassium (Table 1).

The maximum temperature ranged between 31.7°C to 36.1°C, the peak was recorded on the eighth week while the minimum was recorded on the fifth week. The minimum temperature ranged between 21.5°C to 24.4°C and the mean relative humidity ranged between 50.2% to 78% (Table 2).

Table 1. Particle size distribution and chemical properties (0-20 cm)

Soil properties	Values
pH (H ₂ O)	5.71
Organic carbon (g/kg)	9.6
Total Nitrogen (g/kg)	0.72
P (mg/kg)	7
K (cmol/kg)	0.6
Ca (cmol/kg)	7.0
Mg (cmol/kg)	5.79
Mn (mg/kg)	23.6
Fe (mg/kg)	6.0
Cu (mg/kg)	12
Zn (mg/kg)	3.08
Sand (%)	84.5
Clay (%)	9.9
Silt (%)	5.6
Textural Class	Loamy sand

3.1 Mean Height, Collar Diameter and Leaf Production

The effect of watering regime was significant on height of seedlings of *Crescentia cujete* at 5% level of probability (Table 3). The seedlings watered once in five days (W₅) significantly recorded the highest mean value of 20.48 cm, which was not significantly different from the seedlings watered once in four days (W₄) with mean value 20.01 cm, W₆ (19.85 cm) and W₃ (19.99 cm) while seedlings watered once everyday (W₁) had the least mean value of 18.01 cm which was not significantly different from W₂ (18.86 cm) (Table 4).

Table 2. Weekly mean temperatures and relative humidity in the study area

Weeks	Maximum temperature (°C)	Minimum temperature (°C)	Mean relative humidity (%)
1	35.6	21.5	50.2
2	34.3	23.1	56.9
3	34.4	24	69.5
4	35.3	23.5	61
5	31.7	23.9	78
6	32.8	24.3	68.9
7	35	22.2	38.4
8	36.1	24.4	59.8
9	35.4	24.3	64.2
10	35.1	24.2	70.1
11	34.6	24	68.2
12	31.8	23.8	74.9

Source: Forestry Research Institute of Nigeria Meteorology Section

Table 3. Analysis of variance of the growth parameters

Parameters	Source of variation	df	Sum of squares	Mean square	F	Significance
Height	Watering regime	5	25.327	5.065	4.871	0.002*
	Error	30	31.194	1.040		
	Total	35	56.522			
Collar Diameter	Watering regime	5	0.341	0.068	0.622	0.684 ^{ns}
	Error	30	3.292	0.110		
	Total	35	3.633			
Leaf Production	Watering regime	5	59.206	11.841	3.142	0.021*
	Error	30	113.065	3.769		
	Total	35	172.271			

ns – not significant; * - significant ($p \leq 0.05$)

Table 4. Mean values of the growth parameters of *Crescentia cujete* subjected to selected watering regimes

Treatments	Height (cm)	Leaves	Collar diameter (mm)
W ₁	18.01 ^a	15.76 ^a	4.82
W ₂	18.86 ^{ab}	14.56 ^a	4.79
W ₃	19.99 ^{bc}	16.82 ^{ab}	4.65
W ₄	20.01 ^{bc}	14.96 ^a	4.97
W ₅	20.48 ^c	18.42 ^b	4.88
W ₆	19.85 ^{bc}	16.57 ^{ab}	4.81

Means followed by the same superscripts in the same column are not significantly different ($p = .05$)

There was no significant difference in the collar diameter of the seedlings of *Crescentia cujete* at 5% level of probability. However, seedlings watered once in four days (W₄) had the highest mean value of 4.97 mm closely followed by seedlings watered once in five days (W₅) with mean value of 4.88 mm while the seedlings watered once in three days (W₃) had the least mean value of 4.65 mm (Table 4).

In terms of leaf production, there was significant difference among the treatments used at 5% level of probability. The seedlings watered once in five days (W₅) produced the highest number of

leaves with mean value of 18.42 followed by seedlings watered once in three days (W₃) with mean value of 16.82 and the seedlings watered once in two days (W₂) produced the least number of leaves with mean value of 14.56 (Table 4).

3.2 Biomass Accumulation

3.2.1 Leaf biomass accumulation

The result showed that the selected watering regimes had significant effect on the leaf biomass accumulation of *Crescentia cujete* with

Table 5. Mean values of the biomass of sectioned parts of *Crescentia cujete*

Treatments	Leaf dry weight (g)	Stem dry weight (g)	Root dry weight (g)
W ₁	2.52 ^{ab}	1.75 ^b	2.87
W ₂	1.93 ^a	1.18 ^a	2.30
W ₃	2.25 ^{ab}	1.20 ^a	2.37
W ₄	1.82 ^a	1.18 ^a	2.32
W ₅	2.92 ^b	1.80 ^b	3.08
W ₆	2.29 ^{ab}	1.30 ^{ab}	2.18

Means followed by the same superscripts in the same column are not significantly different ($p = .05$)

W₅ recording the highest mean value of 2.29 g and W₄ recording the least mean value which was not significantly different from mean value recorded by W₂ (1.93 g) (Table 5).

3.2.2 Stem biomass accumulation

The result showed that the stem biomass accumulation of *Crescentia cujete* was significantly affected by the selected watering regimes. W₅ had the highest mean value of 1.80 g which is not significantly different from W₁ and W₆ with mean values of 1.75 g and 1.30 g respectively while W₂ and W₄ recorded the least mean value of 1.18 g (Table 5).

3.2.3 Root biomass accumulation

The effect of watering regime on root biomass accumulation of *Crescentia cujete* is not significantly different among the treatments. This implies that irrespective of the selected watering regimes, the root dry weight of *Crescentia cujete* may not differ (Table 5).

4. DISCUSSION

The resultant effect of different watering regimes on the height of *Crescentia cujete* agrees with the findings of [11] who reported that *Parkia biglobosa* subjected to watering once in five days had the best performance in growth. Isah et al. [18] recommended watering once in two to three days for the optimum growth for *Acacia senegal*. Oyun et al. [9] reported that watering twice a week is suitable to achieve the optimum growth of *Acacia senegal* in the nursery.

The seedlings watered once in five days (W₅) produced the highest number of leaves which was not significantly different from the seedlings watered once in three days (W₃) and the seedlings watered once in six days (W₆). [11] observed similar results in studies of different

watering regimes on the growth of *Parkia biglobosa* seedlings where seedlings watered once in three days or once in five days yielded the highest number of leaves. Lisar et al. [19] reported that plants optimize the morphology, physiology and metabolism of their organs and cells in order to maximize productivity under the moisture stress conditions.

Sale [11] also observed that *Parkia biglobosa* seedlings watered once in three days gave the highest dry plant weight. Arndt et al. [20] reported that reduced soil moisture may cause a reduction in root growth and limit nutrient uptake by roots; this is contrary to the result from this study. It has been widely reported that extensive root growth is an adaptive feature under drought stressed conditions [21]. Shoot dry weight increases with increase in soil moisture but under water stress conditions, root growth exhibits better growth than the shoot. This causes the increase in root to shoot ratio [22].

5. CONCLUSION

In this study, the effect of selected watering regimes on *Crescentia cujete* was examined. It is evident that *C. cujete* can optimize water deficit for its growth. For effective water management coupled with optimum growth during the early growth of *C. cujete* seedlings, water can be applied once in four to five days.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Michael A. Trees, shrubs and liannas of West Africa dry zones. Grad Margae Publishers GMBH, MNHN. 2004;191.

2. Gilman EF, Watson DG. Fact sheet florida cooperative extension service. University of Florida; 1993.
3. Ejelonu BC, Lasisi AA, Olaremu AG and Ejelonu OC. The chemical constituents of calabash (*Crescentia cujete*). African Journal of Biotechnology. 2011;10(84): 19631-19636.
4. UNCTAD. Market brief in the European Union for selected natural ingredients derived from native species - *Crescentia cujete* (Calabash, Totumo). United Nations Conference on Trade and Development/ Biotrade Facilitation Programme; 2005.
5. Carter SJ. Tutorial pharmacy. 6th Edn. Pitman Medical Publishing Co. Ltd. Tunbridge Well Kent, England. 1979:29-30.
6. Phaniendra A, Jestadi DB, Periyasamy L. Free radicals: Properties, sources, targets and their implications in various diseases. Indian Journal of Clinical Biochemistry. 2015;30(1):11-26.
7. Ördög V. The agricultural engineering M.Sc curriculum development TÁMOP- 4.1.2-08 / 1 / A- 2009-0010 project; 2011.
8. Price DT, Black TA, Kelliher FM. Effects of sala understorey removal on photosynthesis rate and stomata conductance of young douglas-fir trees. Canadian Journal of Forest Resource. 1986;16:90-97.
9. Oyun MB, Adeduntan SA, Suberu SA. Influence of watering regime and mycorrhizae inoculations on the physiology and early growth of *Acacia senegal* (L.) Willd. African Journal of Plant Science. 2010;4(7):210-216.
10. Levy Y, Krikum J. Effects of irrigation, water and salinity and root-stock on the vertical distribution of vesicular arbuscular mycorrhiza on citrus roots. New Phytology. 1983;15:397-403.
11. Sale FA. Evaluation of watering regimes and different pot sizes in the Growth of *Parkia biglobosa* (Jacq) benth seedlings under nursery condition. European Scientific Journal. 2015;11 (12):313-325.
12. Pallardy SG. Physiology of woody plants. 3rd ed. Academic Press. 2008;95. [ISBN 978-0-12-088765-1]
13. Chaouche K, Neppel L, Dieulin C, Pujol N, Ladouche B, Martin E, et al. Analyses of precipitation, temperature and evapotranspiration in a French mediterranean region in the context of climate change. Comptes Rendus Geoscience. 2010;342(3):234-243.
14. Synder RL, Moratiel R, Zhenwei S, Swelam A, Jomaa I, Shapland T Evapotranspiration response to climate change. 28th International Horticultural Congress on Science and Horticulture for People: International Symposium on clim water: Horticultural use of water in a changing climate. Eds: JE Fernandez, MI Ferrereira. Acts Hort. 2010; 922.
15. Chavarria G, Pessoa dos Santos H. Plant water relations: Absorption, transport and control mechanisms, advances in selected plant physiology aspects, Dr. Giuseppe Montanaro (Ed.); 2012. [ISBN:978-953-51-0557-2] Available:<http://www.intechopen.com/books/advances-in-selected-plantphysiology-aspects/plant-water-relations-absorption-transport-and-control-mechanisms>
16. Shipley B, Vu T. Dry matter content as a measure of dry matter concentration in plants and their parts. New Phytologist. 2002;153:359-364.
17. Steel RG and Torrie JH. Biostatistics: Principles and Procedures. McGraw Hill, Mexico. 1998
18. Isah AD, Bello AG, Maishanu HM and Abdullahi S. Effect of watering regime on the early growth of *Acacia senegal* (linn) willd. Provenances. International Journal of Plant, Animal and Environmental Sciences. 2013;3:52-56.
19. Lisar SYS, Motafakkerazad R, Hossain MM, Rahman IMM. Water stress in plants: Causes, effects and responses. Prof. Ismail Md. Mofizur Rahman (Ed.); 2012. [ISBN: 978-953-307-963-9]
20. Arndt SK, Clifford SC, Wanek W, Jones HG, Popp M. Physiological and morphological adaptation of the fruit tree *Ziziphus rotundifolia* in response to progressive drought stress. Tree Physiology. 2001;21:705-715.
21. Guoxiong C, Krugman T, Fahima T, Korol AB, Nevo E. Comparative

- study on morphological and physiological traits related to drought resistance between xeric and mesic *Hordeum spontaneum* lines in Israel. Barley Genetics Newsletter. 2002;32:22-33.
22. Chiatante D, Di-Iorio A, Sciandra S, Stefania G, Mazzoleni S. Effect of drought and fire on root development in *Quercus pubescens* Willd. and *Fraxinus ornus* L. seedlings. J. Environ. Exp. Bot. 2006; 56:190-197.

© 2019 Aderounmu and Ogidan; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/50674>