



Economics of Wet Seeded Rice as Influenced by Alternate Wetting and Drying Irrigation Regimes and Weed Management

**N. Prathap Reddy^{a*⊙}, G. E. Ch. Vidya Sagar^{a#}, M. Yakadri^{a†}, M. Malla Reddy^{a#},
A. Madhavi^{b‡} and S. Narender Reddy^{c‡}**

^a Department of Agronomy, College of Agriculture, Professor Jayashankar Telangana State Agricultural University, Rajendranagar, Hyderabad -500030, India.

^b AICRP on STCR, ARI, Professor Jayashankar Telangana State Agricultural University, Rajendranagar, Hyderabad -500030, India.

^c Agricultural College, Professor Jayashankar Telangana State Agricultural University Polasa, Jagtial – 505327, India.

Authors' contributions

This work was carried out in collaboration among all authors. Author NPR performed the statistical analysis, wrote the first draft of the manuscript. Authors GECHVS, MY, MMR, AM and SNR designed the study and managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i1131057

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/90236>

Original Research Article

Received 24 May 2022
Accepted 29 July 2022
Published 29 July 2022

ABSTRACT

An experiment was conducted at College Farm, Professor Jayashankar Telangana State Agricultural University, Rajendranagar, Hyderabad during *rabi* 2020-21 and 2021-22. The experiment was laid out in split plot design with two factors i.e., irrigation regimes (I_1 : Farmers practice (continuous flooding of 2-5 cm from 3-4 days after sowing), I_2 : AWDI at 5 cm depletion of ponded water, I_3 : AWDI at 10 cm depletion of ponded water and I_4 : AWDI at 15 cm depletion of ponded water) and weed management practices (W_1 : Control (Unweeded check), W_2 :

[⊙] Ph.D Scholar;

[#] Professor;

[†] Professor and Head;

[‡] Principal scientist, and Head;

[‡] Associate Dean;

*Corresponding author: E-mail: prathapreddynaidi@gmail.com, naiderajureddy27@gmail.com;

Pyrazosulfuron ethyl (10% WP) 20 g ha⁻¹ PE fb Penoxsulam (1.02%) + Cyhalofop butyl (5.1%) 120 g ha⁻¹ PoE, W₃:Pyrazosulfuron ethyl (10% WP) 20 g ha⁻¹ PE fb Penoxsulam (1.02%) + Cyhalofop butyl (5.1%) 120 g ha⁻¹ PoE + Mechanical weeding at 45 DAS and W₄: Weed free (Mechanical weeding at 15, 35 and 55 DAS with line weeding). Among the irrigation regimes, cost of cultivation and gross returns were highest in farmers practice (continuous flooding of 2-5 cm from 3-4 days after sowing) and lowest in AWDI at 15 cm depletion of ponded water. Net returns and B:C ratio was highest in AWDI at 5 cm depletion of ponded water and lowest net returns and B:C ratio was recorded in AWDI at 15 cm depletion of ponded water. Among weed management practices highest cost of cultivation, gross returns were achieved in weed free (mechanical weeding at 15, 35 and 55 DAS with line weeding) and lowest cost of cultivation and gross returns was achieved in Control (unweeded control). With respective to herbicide application highest gross returns, net returns and B:C ratio were achieved in pyrazosulfuron ethyl (10% WP) 20 g ha⁻¹ PE fb penoxsulam (1.02%) + cyhalofop butyl (5.1%) 120 g ha⁻¹ PoE + mechanical weeding at 45 DAS.

Keywords: Wet seeded rice; irrigation regimes; weed management; economics.

1. INTRODUCTION

The most popular method for establishing rice in South East Asian nations, including India, is transplanting. In India, rice is traditionally grown in flooded conditions, however this method requires a lot of water. Water-saving technologies are required in order to continue production even with decreased water use in order to fulfil the increasing demand for food from the population growth. Alternate wetting and drying approach, reduce water input by 30% without lowering yield [1]. Compared to continuous flooding, the AWD greatly reduced the number of irrigations and the amount of irrigation water used.

Heavy weed infestation puts direct seeded rice's viability in jeopardy [2]. In fields that were direct sown, weed growth is typical. Due to the same age and physical characteristics of grass weeds and rice seedlings, rice seedlings in wet seeded conditions experience higher weed growth [3]. Direct seeded rice's productivity may be increased by good weed management practices throughout the first 40 days of crop growth [4]. Herbicides are currently the most crucial weed management tool since they provide fast, efficient, cost-effective, and useful methods of weed control. However, diverse weeds in direct seeded rice could not be adequately controlled by pre or post-emergence herbicide sprays alone [5]. Although manual, mechanical, and chemical weed control methods were effective, lack of labour during peak season and rising labour costs are causing weed control procedures to be delayed and expensive. Therefore, timely weed control is achieved by combining herbicide

applications such as pre and postemergence with mechanical and manual weeding techniques. With these considerations in mind, the current study was conducted to investigate the effects of alternating wetting and drying irrigation regimes, as well as weed management practices, on the economics of wet seeded rice during the *rabi* season.

2. MATERIALS AND METHODS

An experiment was carried out at College Farm, College of Agriculture, Rajendranagar, Professor Jayashankar Telangana State Agricultural University, Rajendranagar, Hyderabad at an altitude of 527 m above mean sea level (MSL), 17° 31' N latitude, 78° 40' E longitude in Southern Telangana zone. The soil of the experimental site was sandy loam in texture with pH 7.9, available nitrogen (251.0 kg ha⁻¹), available phosphorus (42.5 kg ha⁻¹ P₂O₅) and available potassium (364.0 kg ha⁻¹ K₂O).

The experiment was laid out in split plot design with two factors i.e., irrigation regimes (I₁ : Farmers practice (continuous flooding of 2-5 cm from 3-4 days after sowing), I₂: AWDI at 5 cm depletion of ponded water, I₃: AWDI at 10 cm depletion of ponded water and I₄: AWDI at 15 cm depletion of ponded water) and other weed management practices (W₁: Control (Unweeded check), W₂: Pyrazosulfuron ethyl (10% WP) 20 g ha⁻¹ PE fb Penoxsulam (1.02%) + Cyhalofop butyl (5.1%) 120 g ha⁻¹ PoE, W₃:Pyrazosulfuron ethyl (10% WP) 20 g ha⁻¹ PE fb Penoxsulam (1.02%) + Cyhalofop butyl (5.1%) 120 g ha⁻¹ PoE + Mechanical weeding at 45 DAS and W₄: Weed free (Mechanical weeding at 15, 35 and 55 DAS with line weeding). The test variety used for the

study was Jagtial Rice-1 (JGL-24423). Sowing was done by drum seeder by maintaining spacing of 25 cm x 8 cm. The data were analyzed statistically applying analysis of variance technique for split plot design. The significance was tested by 'F' test [6]

2.1 Cost of Cultivation (Rs. ha⁻¹)

Cost of cultivation was calculated by adding the cost of all the inputs viz., seed, fertilizer, herbicides, irrigation water, labour, pesticides etc.

2.2 Gross Returns (Rs. ha⁻¹)

Gross returns were calculated by using the cost of grain and straw in the market at the time of harvesting of the crop and by multiplying them with respective grain and straw yields and expressed as Rs. ha⁻¹

2.3 Net Returns (Rs. ha⁻¹)

Net returns were calculated by subtracting the cost of cultivation from the gross returns and expressed as Rs. ha⁻¹

2.4 Benefit Cost Ratio

Benefit cost ratio was calculated using the formula

$$B - C \text{ Ratio} = \frac{\text{Gross returns (Rs. ha}^{-1}\text{)}}{\text{Cost of cultivation (Rs. ha}^{-1}\text{)}}$$

3. RESULTS AND DISCUSSION

3.1 Cost of Cultivation

Cost of cultivation as influenced by irrigation regimes and weed management practices were presented in Table 1.

The highest cost of cultivation was found with Farmers practice (continuous flooding of 2-5 cm from 3-4 days after sowing) (74370 and 72470.0 Rs ha⁻¹) followed by AWDI at 5 cm depletion of ponded water and AWDI at 10 cm depletion of ponded water. The lowest cost of cultivation resulted in AWDI at 15 cm depletion of ponded water (56170 and 55450 Rs ha⁻¹). This might be due to more number of irrigations applied in farmers practice (continuous flooding of 2-5 cm from 3-4 days after sowing) when compared with

other alternate wetting and drying irrigation regimes. Similar findings were reported by [7]

Cost of cultivation varied with different weed management practices. The highest cost of cultivation was achieved in Weed free (mechanical weeding at 15, 35, and 55 DAS with line weeding) (Rs. 70845 and 70292 Rs ha⁻¹) followed by Pyrazosulfuron ethyl (10% WP) 20 g ha⁻¹ PE fb penoxsulam (1.02%) + cyhalofop butyl (5.1%) 120 g ha⁻¹ PoE + mechanical weeding at 45 DAS) and Pyrazosulfuron ethyl (10% WP) 20 g ha⁻¹ PE fb penoxsulam (1.02%) + cyhalofop butyl (5.1%) 120 g ha⁻¹ PoE when compared with Control (unweeded check) which resulted in the lowest cost of cultivation (55695 and 55142 Rs ha⁻¹). Weed free condition recorded the highest cost of cultivation due to the high requirement of labour for mechanical weeding and line weeding. The lowest cost of cultivation was recorded in unweeded control as there was no usage of labour or herbicide. Similar findings were reported by [8]

3.2 Gross Returns (Rs ha⁻¹)

Gross returns as influenced by irrigation regimes and weed management practices were presented in Table 1.

The highest gross returns in both years of study were found in with Farmers practice (continuous flooding of 2-5 cm from 3-4 days after sowing) (149154 and 156709 Rs ha⁻¹) might be due to higher biological yield in farmers practice (continuous flooding of 2-5 cm from 3-4 days after sowing) and it was on par with by AWDI at 5 cm depletion of ponded water. The lowest gross returns resulted in AWDI at 15 cm depletion of ponded water. (104576 and 110845 Rs ha⁻¹). Similar findings were reported by [7]

Gross returns varied with different weed management practices during both years of study. The highest gross returns were achieved in Weed free (mechanical weeding at 15, 35 and 55 DAS with line weeding) (153021 and 155503 Rs ha⁻¹) and it was on par with Pyrazosulfuron ethyl (10% WP) 20 g ha⁻¹ PE fb penoxsulam (1.02%) + cyhalofop butyl (5.1%) 120 g ha⁻¹ PoE + mechanical weeding at 45 DAS). Application of herbicide alone i.e Pyrazosulfuron ethyl (10% WP) 20 g ha⁻¹ PE fb penoxsulam (1.02%) + cyhalofop butyl (5.1%) 120 g ha⁻¹ PoE resulted in higher gross returns when compared with Control (unweeded check) which resulted in the lowest gross returns. (78974 and 83875 Rs ha⁻¹). Higher

Table. 1. Cost of cultivation and Gross returns of wet seeded rice as influenced by alternate wetting and drying irrigation regimes and weed management practices

Treatments	Cost of cultivation (Rs ha ⁻¹)			Gross returns (Rs ha ⁻¹)		
	20-21	21-22	Mean	20-21	21-22	Mean
Irrigation Regimes (I)						
I ₁ :Farmers practice (continuous flooding of 2-5 cm from 3-4 days after sowing)	74370	72470	73420	149154	156709	152932
I ₂ :AWDI at 5 cm depletion of ponded water	60820	61500	61160	137792	146355	142074
I ₃ : AWDI at 10 cm depletion of ponded water	57820	57550	57685	123615	130864	127239
I ₄ :AWDI at 15 cm depletion of ponded water	56170	55450	55810	104576	110845	107710
S.Em.±	-	-	-	4590	4649	-
CD (p=0.05)	-	-	-	15885	16088	-
Weed Management Practices (W)						
W ₁ :Control (Unweeded check)	55695	55142	55419	78974	83875	82675
W ₂ : Pyrazosulfuron ethyl (10% WP) 20 g ha ⁻¹ PE fb penoxsulam (1.02%) + cyhalofop butyl (5.1%) 120 g ha ⁻¹ PoE	59395	58842	59119	136517	139944	140109
W ₃ :Pyrazosulfuron ethyl (10% WP) 20 g ha ⁻¹ PE fb penoxsulam (1.02%) + cyhalofop butyl (5.1%) 120g ha ⁻¹ PoE + mechanical weeding at 45 DAS	63245	62692	62969	146625	151079	150863
W ₄ :Weed free (mechanical Weeding at 15, 35 and 55 DAS with line weeding)	70845	70292	70569	153021	155503	156308
S.Em.±	-	-	-	2264	2506	-
CD (p=0.05)	-	-	-	6609	7315	-
Interaction						
Weed management practices at same level of irrigation regimes						
S.Em.±	-	-	-	4528.8	5012.7	-
CD (p=0.05)	-	-	-	NS	NS	-
Irrigation regimes at same or different level of weed management						
S.Em.±	-	-	-	6037.7	6360.8	-
CD (p=0.05)	-	-	-	NS	NS	-
General Mean	62295	61742	62018	128784	136193	132488

Table 2. Net returns and B:C ratio of wet seeded rice as influenced by alternate wetting and drying irrigation regimes and weed management practices

Treatments	Net returns (Rs ha ⁻¹)			B: C ratio		
	20-21	21-22	Mean	20-21	21-22	Mean
Irrigation Regimes (I)						
I ₁ :Farmers practice (continuous flooding of 2-5 cm from 3-4 days after sowing)	74784	84239	79512	1.99	2.15	2.07
I ₂ :AWDI at 5 cm depletion of ponded water	76972	84855	80914	2.24	2.36	2.30
I ₃ : AWDI at 10 cm depletion of ponded water	65795	73314	69554	2.12	2.26	2.19
I ₄ :AWDI at 15 cm depletion of ponded water	48406	55395	51900	1.84	1.98	1.91
S.Em.±	4590	4649	-	0.08	0.07	-
CD (p=0.05)	15885	16088	-	0.27	0.25	-
Weed Management Practices (W)						
W ₁ :Control (Unweeded check)	23279	31233	27256	1.42	1.57	1.49
W ₂ : Pyrazosulfuron ethyl (10% WP) 20 g ha ⁻¹ PE fb penoxsulam (1.02%) + cyhalofop butyl(5.1%) 120 g ha ⁻¹ PoE	77122	84858	80990	2.30	2.44	2.37
W ₃ :Pyrazosulfuron ethyl (10% WP) 20 g ha ⁻¹ PE fb penoxsulam (1.02%) + cyhalofop butyl(5.1%) 120g ha ⁻¹ PoE + mechanical weeding at 45 DAS	83380	92409	87894	2.32	2.47	2.39
W ₄ :Weed free (mechanical weeding at 15, 35 and 55 DAS with line weeding)	82176	89303	85739	2.16	2.27	2.21
S.Em.±	2264	2506	-	0.04	0.04	-
CD (p=0.05)	6609	7315	-	0.11	0.12	-
Interactionexcel						
Weed management practices at same level of irrigation regimes						
S.Em.±	4529	5013	-	0.08	0.08	-
CD (p=0.05)	NS	NS	-	NS	NS	-
Irrigation regimes at same or different level of weed management						
S.Em.±	6038	63601	-	0.10	0.10	-
CD (p=0.05)	NS	NS	-	NS	NS	-
General Mean	66489	74451	70470	2.05	2.19	2.12




Fig. 1. Net returns of wet seeded rice as influenced by alternate wetting and drying irrigation regimes and weed management practices

gross returns were achieved in weed free condition followed by W_3 and W_2 due to the higher grain and straw yield in these treatments when compared to unweeded control which resulted in lower gross returns due to lower grain and straw yield. Similar findings were reported by [9]

3.3 Net Returns (Rs ha⁻¹)

Net returns as influenced by irrigation regimes and weed management practices were presented in Table 2 and Fig 1.

The highest net returns during both years of study were found in AWDI at 5 cm depletion of ponded water (76972 and 84855 Rs ha⁻¹) and it was on par with Farmers practice (continuous flooding of 2-5 cm from 3-4 days after sowing) and AWDI at 10 cm depletion of ponded water. This might be due to the lesser cost of cultivation in alternate wetting and drying irrigation at 5 cm depletion of ponded water because of lesser amount of irrigation water and fewer number of irrigations when compared to farmer practice (continuous flooding of 2-5 cm from 3-4 days after sowing). Similar findings were reported by [10]. The lowest net returns resulted in AWDI at 15 cm depletion of ponded water (48406 and 55395 Rs ha⁻¹)

Net returns varied with different weed management practices. Highest net returns was achieved Pyrazosulfuron ethyl (10% WP) 20 g ha⁻¹ PE fb penoxsulam (1.02%) + cyhalofop butyl(5.1%) 120 g ha⁻¹ PoE + mechanical weeding at 45 DAS) (83380 and 92409 Rs ha⁻¹) and it was on par with Weed free (Mechanical weeding at 15, 35, and 55 DAS with line weeding) and Pyrazosulfuron ethyl (10% WP) 20 g ha⁻¹ PE fb penoxsulam (1.02%) + cyhalofop butyl (5.1%) 120 g ha⁻¹ PoE during 2020-21 and during 2021-22, highest net returns was achieved W_3 :Pyrazosulfuron ethyl (10% WP) 20 g ha⁻¹ PE fb penoxsulam (1.02%) + cyhalofop butyl(5.1%) 120 g ha⁻¹ PoE + mechanical weeding at 45 DAS) (92409 Rs ha⁻¹) and it was on par with Weed free (Mechanical weeding at 15, 35, and 55 DAS with line weeding).The lowest net returns were resulted in Control (unweeded check) (23279 and 31233 Rs ha⁻¹). The highest net returns were achieved in W_3 when compared to the weed free condition due to less cost of cultivation incurred in W_3 where herbicide combination + mechanical weeding was carried out. Similar findings were reported by [9] and [8]

3.4 B: C ratio

B:C ratio as influenced by irrigation regimes and weed management practices were presented in Table 2.

The highest B:C ratio during both years of study was found in AWDI at 5 cm depletion of ponded water (2.24 and 2.36) and this was on par with AWDI at 10 cm depletion of ponded water and Farmers practice (continuous flooding of 2-5 cm from 3-4 days after sowing). This might be due to lesser cost of cultivation and more net returns in alternate wetting and drying irrigation at 5 cm depletion of ponded water and alternate wetting and drying irrigation at 10 cm depletion of ponded water because of lesser amount of irrigation water and fewer number of irrigations when compared to Farmer practice (continuous flooding of 2-5 cm from 3-4 days after sowing). Similar findings were reported by [11] and [7] and. The lowest B:C ratio was resulted in I_4 : AWDI at 15 cm depletion of ponded water (1.84 and 1.98)

B:C ratio varied with different weed management practices. Highest B:C ratio during both years of study was achieved in Pyrazosulfuron ethyl (10% WP) 20 g ha⁻¹ PE fb penoxsulam (1.02%) + cyhalofop butyl (5.1%) 120 g ha⁻¹ PoE + mechanical weeding at 45 DAS) (2.32 and 2.47) and this was on par with with Pyrazosulfuron ethyl (10% WP) 20 g ha⁻¹ PE fb penoxsulam (1.02%) + cyhalofop butyl (5.1%) 120 g ha⁻¹ PoE (2.30 and 2.40) followed by Weed free (Mechanical weeding at 15, 35 and 55 DAS with line weeding) (2.16 and 2.30) which might be due to reduced cost of weeding with higher grain and straw yield. Control (unweeded check) resulted in the lowest B:C ratio (1.42 and 1.57) which might be due to lower grain and straw yield in unweeded control. Similar findings were reported by [12].

4. CONCLUSION

Economics of wet seeded rice were significantly influenced by different irrigation regimes and weed management practices. Among irrigation regimes highest gross returns were achieved in farmers practice and it was on par with AWDI at 5 cm depletion of ponded water, highest net returns and B:C ratio was achieved under AWDI at 5 cm depletion of ponded water. Among the weed management practices highest gross returns were noticed under Weed free (mechanical weeding at 15, 35 and 55 DAS with

line weeding). Highest net returns and B:C ratio was noticed in Pyrazosulfuron ethyl (10% WP) 20 g ha⁻¹ PE fb penoxsulam (1.02%) + cyhalofop butyl (5.1%) 120 g ha⁻¹ PoE + mechanical weeding at 45 DAS

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bouman B, Lampayan RM and Tuong TP. Water management in irrigated rice: Coping with water scarcity. International Rice Research Institute, Las Banos, Laguna, Philippines. 2007;53.
2. Mahajan GBS, Chauhan J, Timisina TP and Singh K. Crop performance and water and nitrogen use efficiencies in dry seeded rice in response to irrigation and fertilizer mounts in North West India. Field crops research. 2013;134 :59-70.
3. Ramzan M. Evaluation of various planting methods in rice wheat cropping system. Punjab, Pakistan. Rice Crop Report. 2003;4-5.
4. Maity SK and Mukherjee PK. Integrated weed management in dry direct seeded rainy season rice (*Oryza sativa* L.) Indian Journal of Agronomy. 2008;53(2):116-120.
5. Awan TH, Cruz PC, Chauhan BS. Efficacy and economics of different herbicides, their weed species selectivity, and the productivity of mechanized dry-seeded rice. Crop Protection. 2015;78:239–246.
6. Gomez KA, Gomez AA. Statistical procedure for agriculture research. 2nd Edition. John Willey and Sons, New York. 1984;680.
7. Shabana, Vinod Kumar, Rajan Kumar, Alisha Kumari, Sweeti Kumari. Effect of irrigation management and crop establishment methods on growth, yield, and economics of rice. International Journal of Current Microbiology and Applied Sciences. 2019;8(4):443-451.
8. Mounisha J, Menon MV and Reddy TV. Weed management with ready-mix herbicides in wet seeded rice. Journal of Crop and Weed. 2021;17(1):309-313.
9. Mohanta A, Behura AK. Bio-efficacy of different herbicides on weed dynamics, growth and grain yield of transplanted rice (*Oryza sativa* L.) in Odisha, India. International Journal of Current Microbiology and Applied Science. 2020; 9(4):2258-2563.
10. Rahaman S, Sinha AC. Effect of water regimes and organic sources of nutrients for higher productivity and nitrogen use efficiency of summer rice (*Oryza sativa*). African Journal of Agricultural. 2013;8(48):6189-6195
11. Nayak BR, Pramanik K, Khanda CM, Panigrahy N, Samant PK, Mohapatra, S, Mohanty AK, Dash AK, Panda N, Swain SK. Response of aerobic rice (*Oryza sativa*) to different irrigation regimes and nitrogen levels in western Odisha. Indian Journal of Agronomy, 2016;61(3):321-325.
12. Nagarjun P, Dhanapal GN, Sanjay MT, Yogananda SB, Muthuraju R. Energy budgeting and economics of weed management in dry direct-seeded rice. Indian Journal of Weed Science. 2019; 51:1-5.

© 2022 Reddy et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/90236>