



Phytoconstituents As Bioenhancers: A Review

Chinnarao Piniseti^a and Vimal Arora^{b*}

^a University Institute of Pharma Sciences, Chandigarh University, Mohali, Punjab-140413, India.

^b Department of Pharmacy, University Institute of Pharma Sciences, Chandigarh University, Mohali, Punjab-140413, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i56A33882

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/77570>

Review Article

Received 05 October 2021
Accepted 11 December 2021
Published 13 December 2021

ABSTRACT

Bio-enhancers are the compounds that enhance the bioavailability of active pharmaceutical ingredients without itself having any pharmacological action. Most of them are of natural origin and do not have any side effects. They enhance the bioavailability by influencing variety of mechanisms involved in the drug action like penetration enhancement, improving metabolism, enzyme inhibition, drug targeting etc. Use of these compounds help to reduce the dose frequency which in turn reduces drug retention in turn causing the toxicity and it also helps in developing cost-effective products. Present days these are widely used to enhance the bioavailability of anti-bacterial, anti-viral, antibiotic, anticancer, anti-inflammatory, cardiovascular drugs etc and effective drug targeting. The present review is designed to emphasize the importance of certain phytoconstituents working as bio-enhancers, their classification and different mechanisms of their activity.

Keywords: Bioenhancer; bioavailability; drug targeting; piperine; quercetin; curcumin; emodin.

1. INTRODUCTION

Bio-enhancers are the chemical moieties or majorly phytoconstituents that enhances the plasma concentration and further promote the efficacy of a drug. They don't have any pharmacological activity of their own and get easily eliminated from the body without causing any side effects. They not only increase the bioavailability of active pharmaceutical ingredients like anti-bacterial, anti-fungal, anti-viral, anti-inflammatory, antibiotics, cardiovascular drugs etc. but these can also be used for the effective absorption of nutrients and vitamins. They enhance the bioavailability of drugs using various mechanisms like elevating intestinal absorption, blocking drug metabolising enzymes which further stops liver and intestinal degradation, as well as blocking efflux mechanisms resulting in the blockage of drug elimination by gut as illustrated in the Fig. 1.

There are number of benefits illustrating the benefits of using bioenhancers, including reduction in quantity of dose intake due to enhanced plasma drug concentration, cost reduction due to lower intake of drug, lesser side effects due to lower usage of drug, increased drug tolerance leading to reduction in drug resistance. Better biocompatibility, safety and economy due to their natural origin [1, 2, 3, 4].

2. BIO-ENHANCERS CLASSIFICATION

Based up on the origin of the bioenhancers, these may be classified as of plant origin and animal origin as stated in the Table 1.

2.1 Piperine

Piperine is an alkaloid obtained from the plant *Piper nigrum* and *Piper longum* belonging to the family piperaceae. The raw form of pepper is black coloured, normally used as spice and condiment. Its taste is pungent and it is not soluble in water. Other isomers of piperine like isopiperine, chavicine, isochavicine and piperanine are also available along with piperine in these plants. Piperine has been investigated for various pharmacological properties like neuroprotection, anti-inflammatory, antioxidant and anticancer properties, other than its drug penetration enhancement (1, 2). It has been reported to have bio enhancing effect on vaccines, pyrazinamide, indomethacin, phenytoin, ciprofloxacin ibuprofen and many more [3].

Table 1. Bioenhancers and their origin

Plant origin	Piperine
	Quercetin
	Curcumin
	Allicin
	Glycyrrhizin
	Naringin
	Steviosol
	Niaziridin
	Lysergol
	Emodin
	Simomenine
	Genistein
	5-methoxy hydnocarpin
	Capsaicin
	Capmul
	Camptothecin
Gallic acid	
Ellagic acid	
Ferulic acid	
Animal origin	Cow urine
	Cow urine distillate
	Chitosan

2.2 Quercetin


Quercetin is a natural polyphenolic compound categorized as a flavonoid, having antioxidant potential, It is abundantly available in variety of fruits (apple, grapes, berry and citrus fruits), grains, vegetables (broccoli, onions etc.), wine, coffee, green tea. It is also available in medicinal plants like *Hypericum perforatum*, *Ginkgo biloba* and *Sambucus canadensis*. Piperine has also been investigated for a range of medicinal properties like elevated drug permeability, anti-inflammatory, anti-allergic, antihypertensive, anti-cancer, cardio-protective, anti-viral and antioxidant potential owned via fighting with free radicals. It is also reported to be taken as supplement for energy boost-up [4, 5]. It has been reported to enhance the bioavailability of verapamil, diltiazem, paclitaxel, digoxin, tamoxifen etc. [3].

2.3 Curcumin

Curcumin is a brilliant yellow to orangish yellow coloured compound obtained from *Curcuma longa* (turmeric) belonging to the family Zingiberaceae. Turmeric is known as the wonder drug in the Indian traditional medicine system, as it is used for the treatment of massive number of ailments. It is the major compound of the turmeric extract (Curcumin - 60–70%,

demethoxycurcumin - 20–27%, bisdemethoxycurcumin - 10–15%) and widely used in herbal medicines, cosmetics, and food industry as a colouring and flavouring agent.

Curcumin is a natural phenolic compound responsible for yellow colour of turmeric. It is used as antiviral, antibiotic and anticancer agent [6, 7].



(2*E*,4*E*)-5-(benzo[*d*][1,3]dioxol-5-yl)-1-(piperidin-1-yl)penta-2,4-dien-1-one

Chemical Formula: C₁₇H₁₉NO₃


Exact Mass: 285.14

Molecular Weight: 285.34

m/z: 285.14 (100.0%), 286.14 (18.7%), 287.14 (2.3%)

Elemental Analysis: C, 71.56; H, 6.71; N, 4.91; O, 16.82

Fig. 1. Chemical structure of piperine



2-(3,4-dihydroxyphenyl)-3,5,7-trihydroxy-4*H*-chromen-4-one

Chemical Formula: C₁₅H₁₀O₇


Exact Mass: 302.04

Molecular Weight: 302.24

m/z: 302.04 (100.0%), 303.05 (16.6%), 304.05 (2.7%)

Elemental Analysis: C, 59.61; H, 3.33; O, 37.06

Fig. 2. Chemical structure of quercetin



(1*E*,6*E*)-1,7-bis(4-hydroxy-3-methoxyphenyl)hepta-1,6-diene-3,5-dione

Chemical Formula: C₂₁H₂₀O₆

Exact Mass: 368.13

Molecular Weight: 368.38

m/z: 368.13 (100.0%), 369.13 (23.2%), 370.13 (3.7%)

Elemental Analysis: C, 68.47; H, 5.47; O, 26.06

Fig. 3. Chemical structure of curcumin

2.4 Allicin

Allicin is a sulphurous organic chemical compound obtained from the plant *Alium sativum* (garlic) belonging to the family alliaceae. It is light yellow coloured oily substance that is accountable for distinctive odour of garlic. Biologically it is used in the treatment of septicaemia occurring due to virus and fungus and as an antioxidant [8, 9] It has been found to have bio-enhancing effect on Amphotericin-B and many other drugs [3].

2.5 Ginger

It is the rhizome part of the plant *Zingiber officinale* belonging to family Zingiberaceae. The rhizomes of ginger are popularly used as flavouring agent in food, whereas the dried ginger is used a spice and condiment. The colour of the rhizome varies depending on the variety from yellowish white to light brown with some purplish patches. Traditionally, ginger is provided with lots of health benefits like preventing motion sickness, headache, nausea, and vomiting. It also shows anti-inflammatory potential due to presence of gingerol, anti-cancer mainly towards colorectal cancer and it also helps in killing

tumour cells in case of ovarian cancer. It is reported to be acting as an immuno-booster in various research studies and ancient literature [10,11,12,13,14]. It is found to be possessing bioenhancing effect on Azithromycin, Erythromycin, Cephalexin, Cefadroxil, Amoxycillin, Cloxacillin etc. [3].

2.6 Glycyrrhizin

Glycyrrhizin is the sweetening constituent derived from the root of plant *Glycyrrhiza glabra* (mulethi). It is extensively used as the sweetening agent in various pharmaceutical and nutraceutical preparations. Along with it, the use of this herb is also reported to be as flavouring agent in food and beverage industry, as conditioner as well as humectant in cosmetics for preserving moisture content. In the traditional system of medicine, it is reported to be very useful in the treatment of cough, sore throat, severe hepatitis condition at predefined amounts. An interesting fact about this drug is that at high doses it may lead to high blood pressure and electrolyte imbalance [15,16]. It is reported to be a potential source of increasing bioavailability of rifampicin, tetracycline, nalidixic acid, ampicillin some other antibiotics [3].




Fig. 4. Chemical structure of allicin




Fig. 5. Chemical structure of gingerol




Fig. 6. Chemical structure of glycyrrhizin




Fig. 7. Chemical structure of naringin

2.7 Naringin

It is a naturally occurring flavonoid available in citrus fruits (bergamot, orange, lemon,

mandarin), grapes and is responsible for the sour taste of these fruits. It possesses powerful anti-inflammatory, antioxidant property and is used in treatment of obesity, high blood sugar and high

blood pressure. This is also used in food industry as flavouring agent and to improve the texture of the food. It is also used prevent bitterness in case of canned citrus food. In pharmaceutical industry it is explored for its bioenhancing potential [17, 18]. It is reported to enhance the bioavailability of Verapamil, Tamoxifen, Quinine, Clopidogrel, nimodipine, diltiazem etc. [3].

2.8 Cuminum

The seeds of the plant *cuminum cyminum* belonging to the family apiaceae are used extensively as carminative in the traditional system of medicine in India. These are dark brown coloured seeds with aromatic flavour (cuminaldehyde, cymene, terpinoids) and used as spice. It helps in the treatment of belching, gastric refreshment, removal of helminthes from gastrointestinal tract. Oil (essential oil) derived from the seeds of cuminum is used in cosmetics and is reported to be responsible for its bioenhancer potential [19, 20]. It is reported to improve the bioavailability of Ketoconazole, Erythromycin, Fluorouracil, Amoxicillin, Fluconazole, Cephalixin, Zidovudine etc. [3].


2.9 Carum carvi

The seeds obtained from the plant of *Carum carvi* belonging to the family apiaceae is commonly known as caraway. It is having similarities with the seeds of plant *cuminum cyminum* in terms of their shape and size. The aroma of the seeds is due to carvone, limonene and anethole. Caraway seeds are used as carminative and it is reported to be very useful in providing an effective relief from burping, loosening the bowel for easy defecation, and it also helps in elimination of excess body fluids via diuresis. They improved plasma concentration of antiviral, antifungal, antibiotics and chemotherapy drugs have been reported in the presence of caraway [21, 22].

2.10 Stevia

Stevia is known for its sweetening potential and is reported to be many times sweeter than sugar. It is obtained from the leaves of the plant *Stevia rebaudiana*. The sweetness is owed to the presence of two glycosides stevioside and rebaudioside. It is explored to be used as an additive, food supplement and an alternate to sugar as well as synthetic sweeteners, especially for the benefit of the diabetic patients and elderly

people which help in low intake calories [23, 24, 25, 27]. It has also been explored for its bioenhancing activity in the last few decades, demonstrating it as its active potential.



2-methyl-5-(prop-1-en-2-yl)cyclohex-2-enone

Chemical Formula: C₁₀H₁₄O

Exact Mass: 150.10

Molecular Weight: 150.22

m/z: 150.10 (100.0%), 151.11 (11.0%)

Elemental Analysis: C, 79.96; H, 9.39; O, 10.65


Fig. 8. Chemical structure of carvone

2.11 Niaziridin

Niaziridin is a glycoside extracted from the leaves, pods and bark of the plant *Moringa oleifera*. Medicinally it is well known to be used in the treatment of arthritic pain, and hyperlipidaemia, along with its anti-teratogenic potential to provide protection to the embryo and foetus. As per the Indian traditional system of medicine it is also used for its potential activities such as hepatoprotection, anti-fertility, anti-microbial, anti-fungal, anti-cancer, anti-inflammatory, spasmolytic, anti-ulcer and antioxidant profile [28, 29]. It is found to have bio-enhancing effect on anti-fungal drugs etc. [3].

2.12 Lysergol

It is a clavinet alkaloid belonging to the class of ergoline and is obtained from the plant *Rivea corymbosa* belonging to family convolvulaceae. It is also reported to be present in very small amount in the claviceps group of fungi, and the seeds of the plant *Ipomoea muricata*. Medicinally it is reported to have neuroleptic properties and thus is used for treating psychosis, especially post-surgical psychosis. It is also reported to be used as a psychotropic analgesic, hypotensive and uterus as well as intestine stimulant. But is found to have many neurological side effects like, hallucinations, seizures, delirium, burning sensations and gangrene [30, 31]. The limitations of its side effect and its potential to enhance the bioavailability of the drugs has given a new direction to explore this active alkaloid.



Chemical Formula: $C_{39}H_{64}O_{18}$


Exact Mass: 820.41

Molecular Weight: 820.92

m/z: 820.41 (100.0%), 821.41 (42.2%), 822.42 (8.7%), 822.41 (3.7%), 823.42 (1.6%)

Elemental Analysis: C, 57.06; H, 7.86; O, 35.08

Fig. 9. Chemical structure of stevioside



2-(4-((3,4,5-trihydroxy-6-methyltetrahydro-2H-pyran-2-yl)oxy)phenyl)acetonitrile

Chemical Formula: $C_{14}H_{17}NO_5$


Exact Mass: 279.11

Molecular Weight: 279.29

m/z: 279.11 (100.0%), 280.11 (15.7%), 281.12 (1.1%), 281.11 (1.1%)

Elemental Analysis: C, 60.21; H, 6.14; N, 5.02; O, 28.64

Fig. 10. Chemical structure of niaziridin



((6a*R*,9*R*)-7-methyl-4,6,6a,7,8,9-hexahydroindolo[4,3-*fg*]quinolin-9-yl)methanol

Chemical Formula: $C_{16}H_{18}N_2O$

Exact Mass: 254.14

Molecular Weight: 254.33

m/z: 254.14 (100.0%), 255.15 (17.6%), 256.15 (1.7%)

Elemental Analysis: C, 75.56; H, 7.13; N, 11.01; O, 6.29

Fig. 11. Chemical structure of lysergol

2.13 Aloe vera


Aloe vera is a plant with succulent and thick leaves belonging to the family asphodelaceae and reported to be used traditionally for the treatment of number of ailments. The leaves of this plant contains gel and mainly comprising of constituents like anthraquinones (emodin, lectin), mannans, polymannans and, anthrones. The gel separated from the leaves is used in the preparation of topical products for skin care, burns and cosmetic preparations [32, 33, 34, 35, 36]. The juice of this plant is also reported to be used for weight loss and digestant property along with its use as a food in desert areas of India. It is also used for its emollient and skin hydration property and is thus applied directly on the skin surface for its protective action. It is also used as a thickening and protective agent in variety of cosmetic products. It has also been envisaged for its potential to increase the absorption of vitamins and insulin [3].

2.14 Sinomenine

An alkaloid obtained from the roots of climber *Sinomenium acutum* and also known by the name of cocculine. It is used as an herbal medicine for the treatment of joint pain and inflammation [37]. It is reported to be having bioenhancing effect on digoxin, quinidine, paeniflorin, verapamil etc. [3].

2.15 Genistein

Isoflavone derived from the plant *Genista tinctoria*, *Glycine max* and *Pueraria lobata*; also called as growth inhibitor and phytoestrogen. Biologically found to have antioxidant, anthelmintic, anti-cancer and atherosclerosis potential [38, 39, 40, 41]. In various research studies it has been proven that genistein can enhance the bioavailability of two anticancer drugs namely paclitaxel and vinblastine. [3].



1,3,8-trihydroxy-6-methylantracene-9,10-dione

Chemical Formula: $C_{15}H_{10}O_5$


Exact Mass: 270.05

Molecular Weight: 270.24

m/z: 270.05 (100.0%), 271.06 (16.5%), 272.06 (2.3%)

Elemental Analysis: C, 66.67; H, 3.73; O, 29.60

Fig. 12. Chemical structure of emodin



(4b*R*,8a*S*,9*S*)-4-hydroxy-3,7-dimethoxy-11-methyl-9,10-dihydro-5*H*-9,4b-(epiminoethano)phenanthren-6(8a*H*)-one

Chemical Formula: $C_{19}H_{23}NO_4$


Exact Mass: 329.16

Molecular Weight: 329.39

m/z: 329.16 (100.0%), 330.17 (21.0%), 331.17 (2.9%)

Elemental Analysis: C, 69.28; H, 7.04; N, 4.25; O, 19.43

Fig. 13. Chemical structure of sinomenine



5,7-dihydroxy-3-(4-hydroxyphenyl)-4*H*-chromen-4-one

Chemical Formula: C₁₅H₁₀O₅


Exact Mass: 270.05

Molecular Weight: 270.24

m/z: 270.05 (100.0%), 271.06 (16.5%), 272.06 (2.3%)

Elemental Analysis: C, 66.67; H, 3.73; O, 29.60

Fig. 14. Chemical structure of genistein



5,7-dihydroxy-2-((2*R*,3*R*)-3-(4-hydroxy-3,5-dimethoxyphenyl)-2-(hydroxymethyl)-2,3-dihydrobenzo[*b*][1,4]dioxin-6-yl)-4*H*-chromen-4-one

Chemical Formula: C₂₆H₂₂O₁₀

Exact Mass: 494.12

Molecular Weight: 494.45

m/z: 494.12 (100.0%), 495.12 (28.1%), 496.13 (6.0%)

Elemental Analysis: C, 63.16; H, 4.48; O, 32.36

Fig. 15. Chemical structure of 5-Methoxyhydnicarpin

2.16 5-Methoxyhydnicarpin (5-MHC)

5-MHC is obtained from the plant *Hydnocarpus wightianus*, as well as berberis species; chemically it is an amphipathic weak acid. It is one of the active components of chaulmoogra oil, used for the treatment of leprosy traditionally. It is found to have anti-cancer activity and enhances the antimicrobial activity of berberine [42, 43, 44].

2.17 *Ammannia multiflora*

Herbaceous plant often called as many flower ammannia belonging to family Lythraceae. Used commercially as decorative purpose and in the preparation of baked food. It has been reported that it contains ammaniol, an alcoholic compound to have bio-enhancing potential. Found to have

bio-enhancing effect on anti-microbial agents and few antibiotics like nalidixic acid [45, 46].

2.18 Capsaicin


Capsaicin (8-methyl-N-vanillyl-6-nonenamide) is one of the active phytoconstituents found in the plants of genus capsicum i.e., Chilli peppers. Capsaicin and its chemical constituents together are called as capsaicinoids; found to be causing irritation and burning sensation when encounters the skin surface. It is reported to be used in the formulations used topically for pain relief caused by aches, arthritis, back pain, etc. Also found to reduce peripheral neuropathy, post-herpetic neuralgia, diabetic neuropathy, and cardiovascular diseases. It has been reported to

be showing bio-enhancing effect on theophylline and fexofenadine [47, 48, 49, 50].

2.19 Capmul

Capmul is a diversified group of multifunctional lipids, majorly mono and diglycerides, obtained

from vegetable oil fatty acids and esterification of propylene glycol and glycerine with certain vegetable oil fatty acids. It is used as an emulsifying agent and widely used to enhance bioavailability, solubility, and penetration [51]. One of the major components of capmul is monocaprylin.



(*E*)-*N*-(4-hydroxy-3-methoxybenzyl)-8-methylnon-6-enamide

Chemical Formula: $C_{18}H_{27}NO_3$


Exact Mass: 305.20

Molecular Weight: 305.41

m/z: 305.20 (100.0%), 306.20 (20.0%), 307.21 (1.9%)

Elemental Analysis: C, 70.79; H, 8.91; N, 4.59; O, 15.72

Fig. 16. Chemical structure of ammaniol



(*E*)-*N*-(4-hydroxy-3-methoxybenzyl)-8-methylnon-6-enamide

Chemical Formula: $C_{18}H_{27}NO_3$


Exact Mass: 305.20

Molecular Weight: 305.41

m/z: 305.20 (100.0%), 306.20 (20.0%), 307.21 (1.9%)

Elemental Analysis: C, 70.79; H, 8.91; N, 4.59; O, 15.72

Fig. 17. Chemical structure of capsaicin



Chemical Formula: $C_{11}H_{22}O_4$

Exact Mass: 218.15

Molecular Weight: 218.29


m/z: 218.15 (100.0%), 219.16 (11.9%)

Elemental Analysis: C, 60.52; H, 10.16; O, 29.32

Fig. 18. Chemical structure of monocaprylin

2.20 Peppermint Oil

Peppermint oil is the volatile oil extracted from the plant belonging to the mint family, especially *Mentha piperita*. The two important chemical constituents of peppermint oil are menthone, and menthol which are used as insect repellents and are used traditionally for treatment of variety of ailments. Medicinally menthol is used in the treatment of irritable bowel syndrome, neuropathy, muscle pain, itching, decongestion, and to reduce heart burns. (52, 53, 54, 55, 56, 57) Menthol is a cyclic alcohol and more specially is classified as monoterpenoid, which can also be synthesized synthetically in the laboratory.



Chemical Formula: C₁₀H₂₀O


Exact Mass: 156.15

Molecular Weight: 156.27

m/z: 156.15 (100.0%), 157.15 (10.8%)

Elemental Analysis: C, 76.86; H, 12.90; O, 10.24

Fig. 19. Chemical structure of menthol



3,4,5-trihydroxybenzoic acid

Chemical Formula: C₇H₆O₅

Exact Mass: 170.02

Molecular Weight: 170.12

m/z: 170.02 (100.0%), 171.02 (7.6%), 172.03 (1.3%)

Elemental Analysis: C, 49.42; H, 3.55; O, 47.02

Fig. 20. Chemical structure of gallic acid

2.21 Gallic Acid

Gallic acid is a solid phenolic acid commonly found in variety of plants like sumac, witch, tea leaves, hazel, gallnuts, oak bark etc. It is synthesized by the process of hydrolysis of


tannins and the products are called gallotannins and ellagitannins. It is reported to be used as antioxidant, anti-hyperlipidaemic and commercially as writing ink, along with its utilization in the process of tanning process and manufacturing of paper. It has also shown bio-enhancing effect on anti-viral drugs, diltiazem, metoprolol etc. [58, 59, 60].

2.22 Ellagic Acid

Ellagic acid is a phenolic compound found in vegetables and fruits like walnuts, pecans, cranberries, raspberries, strawberries, chestnuts, grapes, peaches and pomegranates. Used as antioxidant and anti-proliferative agent. Medicinally used as dietary additive, anti-cancer agent and in curing heart diseases [61, 62, 63, 64, 65, 66].

2.23 Ferulic Acid

Ferulic acid is an organic phenol compound found in plant cell walls mainly fennel ferula communis. It is found in vegetables mainly bamboo shoot, popcorn, flaxseed and barley grain, bran, wheat. It is used in the treatment of osteoporosis, Alzheimer's disease, atherosclerosis, cancer, hyperlipidaemic condition, diabetes, menopause problems, and antioxidant and in skin treatment [67, 68, 69, 70, 71, 72].



Chemical Formula: C₁₄H₆O₈

Exact Mass: 302.01

Molecular Weight: 302.19


m/z: 302.01 (100.0%), 303.01 (15.1%), 304.01 (1.6%), 304.01 (1.1%)

Elemental Analysis: C, 55.64; H, 2.00; O, 42.35

Fig. 21. Chemical structure of gallic acid


2.24 Sodium Caprate

Sodium caprate is a natural compound known to be used as one of the efficient intestinal permeation enhancers, majorly used in the delivery of macromolecules. It is the sodium salt of an aliphatic saturated 10-carbon medium chain fatty acid i.e., capric acid. It is also enlisted as the FDA approved food additive [73, 74].



Chemical Formula: C₁₀H₁₀O₄
Exact Mass: 194.06
Molecular Weight: 194.19
m/z: 194.06 (100.0%), 195.06 (10.8%)
Elemental Analysis: C, 61.85; H, 5.19; O, 32.96

Fig. 22. Chemical structure of ferulic acid



sodium decanoate
Chemical Formula: C₂₀H₃₈Na₂O₄
Exact Mass: 388.26
Molecular Weight: 388.49
m/z: 388.26 (100.0%), 389.26 (22.2%), 390.26 (3.1%)
Elemental Analysis: C, 61.83; H, 9.86; Na, 11.84; O, 16.47

Fig. 23. Chemical structure of sodium caprate




Fig. 24. Mechanism of action of bioenhancers

3. MECHANISM OF ACTION OF BIOENHANCERS

There are variety of mechanisms reported in several research studies regarding the permeation enhancing potential of different bioenhancers. Few of these mechanisms are illustrated here under in the figure:

4. CONCLUSION

It is evident from number of research studies that most of the newly developed pharmaceutically active compounds, especially the biomacromolecules like proteins and peptides have either lower bioavailability or lower biostability. In this direction the bioenhancers are the right choice of compounds to overcome these limitations. Even though the surfactants can be an effective solution but naturally occurring bioenhancers are preferred over these because of safety, biocompatibility, low cost and abundant availability. This results in reduction of cost, toxic side effects, and lower drug dose which in turn assure better therapeutic profile. Therefore, this article emphasized on the significance of naturally occurring bioenhancers and their aiming to highlight the need of exhaustive exploration of natural resources as well processing of established naturally occurring bioenhancers to design innovative and efficient drug delivery system. So further research studies must be continued in developing new formulations using these bio-enhancers to target efficient and affordable drug delivery system availability to the economically weaker sections of the society.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kunisuke Izawa, Motonaka Kuroda. Chemical ecology. Comprehensive Natural Products II. 2020; 4:631-671.

2. Elroy Saldanha, Manjeshwar Shrinath Baliga. Health effects of various dietary agents and phytochemicals. Therapy of Acute Pancreatitis” in Therapeutic, Probiotic and Unconventional Foods, Science Direct. 2018;chapter-17:303-314.

3. Sindhoora D, Ananya Bhattacharjee, Shabaraya AR, Gurpreet Kaur Randhawa, Jagdev Singh Kullar, Rajkumar. Bioenhancers from mother nature and their applicability in modern medicine- a comprehensive review, Int J Appl Basic Med Res. Jan 2011;1:5-10.

4. Ryan Raman MS. RD, Kathy W. Warwick. R.D., CDE, What is Quercetin – Benefits, Foods, Dosage, and Side Effects. Available:<https://www.healthline.com/nutrition/quercetin> accessed on 10 Jan, 2020.

5. Yao Li, Jiaying Yao, Chunyan Han, Jiaxin Yang, Maria Tabassum Chaudhry, Shengnan Wang, Hongnan Liu, Yulong Yin. Review on quercetin, inflammation and immunity. Nutrients. 15 March 20f16;8(3).

6. Majeed S, The State of the Curcumin Market, Natural Products Insider; 28 December 2015. Available:<https://www.naturalproductsinsider.com/herbs-botanicals/state-curcumin-market> Accessed on 10 Jan 2021.

7. Nelson KM, Dahlin JL, Bisson J, Graham J, Pauli GF, Walters MA. The essential medicinal chemistry of curcumin. Journal of Medicinal Chemistry. 11 Jan 2017;1620-1637.

8. Eric Block. The chemistry of garlic and onions. Scientific American. April 1985;252(3):114-9.

9. Cavallito CJ, Bailey JH. Allicin, the antibacterial principle of *Allium sativum* - Isolation, Physical Properties and Antibacterial Action. Journal of the American Chemical Society. November 1; 1944:1950-1951.

10. Ginger, NCCIH Herbs at a Glance. US NCCIH; 1 September 2016. Available:<https://www.nccih.nih.gov/health/ginger> Accessed on 15 Jan 2021.

11. Viestad A. Where flavour was born: Recipes and culinary travels along the Indian Ocean Spice Route. Chronicle Books San Francisco. 2007;210-235.

12. Ross M, Pawley A, Osmond M. The lexicon of Proto Oceanic: The culture and environment of ancestral Oceanic society.

- Oceanic Linguistics. January 2014; 3:524-527.
13. Robert B, Trussel S. The Austronesian comparative dictionary: A work in progress. Ocean. Linguist. December 2013;52:493-523.
 14. George Mateljan. World's healthiest foods. GMF Publishing; 2nd edition. 2015;180-187.
 15. Licorice and licorice derivatives, US Food and Drug Administration, Code of Federal Regulations, CFR-2000, title 21: 3:sec 184-1408. 507-508.
 16. Glycyrrhizin, National Library of Medicine, National Center for Biotechnology Information.
Available:<https://pubchem.ncbi.nlm.nih.gov/compound/Glycyrrhizin>
Accessed on 15 Jan 2021.
 17. Ashrafal Alam M, Nasratsubhan, Mahbuburrahman M, Shaikh J. Uddin, Hasan M. Reza, Satyajit D. Sarker. Effect of citrus flavonoids, naringin and naringenin, on metabolic syndrome and their mechanisms of action. Advances in Nutrition. July 2014;5:404-417.
 18. Maria H. Ribeiro. Naringinases: Occurrence, characteristics, and applications. Applied Microbiology and Biotechnology. 2011;90(6):1883-1895.
 19. Ratan deep Singh, Bhavsar Shailesh, Sarita Devi, Jatin Patel. Indian herbal bioenhancers: A review. Pharmacognosy Reviews. January 2009;80-82.
 20. Kesarwani K, Gupta R, Mukerjee A. Bioavailability enhancers of herbal origin: An overview. Asian Pac J Trop Biomed. 2013;3(4):253-266.
 21. USDA Plants Database. *Carum carvi*.
Available:<https://plants.sc.egov.usda.gov/home/plantProfile?symbol=CACA19>
Accessed on 16 Jan 2021.
 22. Meeran Mohamed Fizur, Javed Hayate, Al Tae Hasan, Azimullah Sheikh, Ojha Shreesh K. Pharmacological properties and molecular mechanisms of thymol: Prospects for its therapeutic potential and pharmaceutical development. Frontiers in Pharmacology. 2017;8:380.
 23. Bhatt et al. Evaluation of cardioprotection and bio-efficacy enhancement of stevioside and diltiazem in rats. Future Journal of Pharmaceutical Sciences. 2020;6:34, 13.
 24. Saikat Dewanjee et al. Plant-based antidiabetic nanoformulations: The emerging paradigm for effective therapy. Int. J. Mol. Sci. 2020;21:2217.
 25. Smruti Ranjan Pradhan. A comparative study on anti-microbial activity exhibited by *Stevia rebaudiana* l. Bertoni in Methanol and Chloroform Extract, International Journal of Science and Research. 2018;7(5):904-909.
 26. Goyal SK, Samsher; Goyal RK. Stevia (*Stevia rebaudiana*) a bio-sweetener: A review. International Journal Food Science Nutr. February 2010;1-10.
 27. Momtazi-Borojeni AA, Esmaeili SA, Abdollahi E, Sahebkar A. A Review on the pharmacology and toxicology of steviol glycosides extracted from *Stevia rebaudiana*. Curr Pharm Des. 2017; 23(11):1616-1622.
 28. Ghanshyam B. Dudhatra, Shailesh K. Mody, Madhavi M. Awale, Hitesh B. Patel, Chirag M. Modi, Avinash Kumar, Divyesh R. Kamani, Bhavesh N. Chauhan. A comprehensive review on pharmacotherapeutics of herbal bio enhancers. The Scientific World Journal. August 2012; 2012:637953.
 29. Shanker K, Gupta MM, Srivastava SK, Bawankule DU, Pal A, Khanuja SPS. Determination of bioactive nitrile glycoside(s) in drumstick (*Moringa oleifera*) by reverse phase HPLC. Food Chemistry. 2007;376-382.
 30. Yuan, Haosen, Guo, Zhixian, Luo, Tuoping. Synthesis of (+)-lysergol and its analogues to assess serotonin receptor activity. Org. Lett. Jan 20, 2017;624-627.
 31. Lysergol, National Library of Medicine, National Center for Biotechnology Information.
Available:<https://pubchem.ncbi.nlm.nih.gov/compound/14987>
Accessed on 12 Jan 2021.
 32. Dr. Sanjay Kholiya, Dr. Jyoti Gangwal, Dr. Rajendra Prasad Sharma. Composition and applications of aloe vera gel. International Journal of Trend in Scientific Research and Development. 2020;4 (3):1033-1035.

33. Yates A. Yates garden guide, Harper Collins Australia. 2015;44th Edition:322-335.
34. King GK, Yates KM, Greenlee PG, Pierce KR, Ford CR, McAnalley BH, Tizard IR. The effect of Acemannan Immunostimulant in combination with surgery and radiation therapy on spontaneous canine and feline fibrosarcomas. *J Am Anim Hosp Assoc.* 1995 Sep-Oct;31(5): 439-47.
35. Eshun K, He Q. Aloe vera: a valuable ingredient for the food, pharmaceutical and cosmetic industries - A review. *Critical Reviews in Food Science and Nutrition.* 2004;91-96.
36. Aloe vera, National Centre for Complementary and Integrative Health, US National Institutes of Health. Available:<https://www.nccih.nih.gov/health/aloe-vera> Accessed on 12 Jan 2021.
37. Zhao ZZ, Liang ZT, Zhou H, Jiang ZH, Liu ZQ, Wong YF, Xu HX, Liu L. Quantification of sinomenine in caulis sinomenii collected from different growing regions and wholesale herbal markets by a modified HPLC method. *Biol Pharm Bull.* 2005 Jan;28(1):105-9.
38. Sail, Vibhavari; Hadden, M. Kyle, Desai, Manoj C. (ed.). Chapter Eighteen - notch pathway modulators as anticancer chemotherapeutics. *Annual Reports in Medicinal Chemistry, Annual Reports in Medicinal Chemistry, Academic Press;* 2012.
39. Rao HSP, Reddy KS. Isoflavones from *Flemingia vestita*. *Fitoterapia.* 1991;62(5):458.
40. Si Hongwei, Liu Dongmin, Si Hongwei, Liu Dongmin. Phytochemical genistein in the regulation of vascular function: New Insights Current Medicinal Chemistry, February 2007;2581-2589.
41. Morito Keiko, Hirose Toshiharu, Kinjo Junei, Hirakawa, Tomoki, Okawa Masafumi, Nohara Toshihiro, Ogawa Sumito, Inou, Satoshi, Muramatsu Masami, Masamune Yukito. Interaction of phytoestrogens with estrogen receptors α and β . *Biological & Pharmaceutical Bulletin.* 2001;24: 351-356.
42. Stermitz FR, Lorenz P, Tawara JN, Zenewicz LA, Lewis K. Synergy in a medicinal plant: antimicrobial action of berberine potentiated by 5'-methoxyhydrnocarpin, a multidrug pump inhibitor. *Proceedings of the National Academy of Sciences.* February 15, 2000;1433-1437.
43. Williams, Cheryll. *Medicinal plants in Australia: An antipodean apothecary.* Rosenberg Publishing. 2013;4:297-319.
44. Xiao Yan Xu, Hui Ruiwang, Shu Li Guo, Bo li, 5-Methoxyhydrnocarpin shows selective anticancer effects and induces apoptosis in THP-1 human leukemia cancer cells via mitochondrial disruption, suppression of cell migration and invasion and cell cycle arrest. *Bangladesh Journal of Pharmacology.* June 2016; 11(3):652-60.
45. Upadhyay HC, Dwivedi GR, Darokar MP, Chaturvedi V, Srivastava SK. Bioenhancing and antimycobacterial agents from *Ammannia multiflora*. *Planta Med.* 2012 Jan;78(1):79-81.
46. Armel Jackson Seukep, Victor Kuete, Lutfun Nahar, Satyajit D. Sarker, Mingquan Guo, Plant-derived secondary metabolites as the main source of efflux pump inhibitors and methods for identification. *Journal of Pharmaceutical Analysis.* 2020;10(4):277-290.
47. Peppin JF, Pappagallo M. Capsaicinoids in the treatment of neuropathic pain: A review. *Ther Adv Neurol Disord.* 2014;7(1):22-32.
48. Qin Y, Ran L, Wang J, et al. Capsaicin supplementation improved risk factors of coronary heart disease in individuals with low HDL-C Levels., *Nutrients.* 2017;9(9):1037.
49. Fattori V, Hohmann MS, Rossaneis AC, Pinho Ribeiro FA, Verri WA. Capsaicin: Current Understanding of Its Mechanisms and Therapy of Pain and Other Pre-Clinical and Clinical Uses, *Molecules.* July 2016;21(7):844.
50. Derry S, Sven-Rice A, Cole P, Tan T, Moore RA. Topical capsaicin (high concentration) for chronic neuropathic pain in adults. *Cochrane Database Syst Rev.* 2013 Feb;28(2):CD007393.
51. Hetal P. Thakkar, Amit Khunt, Rahul D. Dhande, Arpita A. Patel. Formulation and evaluation of Itraconazole nanoemulsion for enhanced oral bioavailability. *Journal of Microencapsulation.* 2015;32(6):559-569.

52. Chumpitazi BP, Kearns GL, Shulman RJ. Review article: the physiological effects and safety of peppermint oil and its efficacy in irritable bowel syndrome and other functional disorders, *Aliment Pharmacol Ther.* 2018;47(6):738-752.
53. Kumar Sarita, Wahab Naim, Warikoo Radhika, Bioefficacy of *Mentha piperita* essential oil against dengue fever mosquito *Aedes aegypti*. *Asian Pacific Journal of Tropical Biomedicine.* April 2011; 1:85-88.
54. Babar Ali, Naser Ali Al-Wabel, Saiba Shams, Aftab Ahamad, Shah Alam Khan, Firoz Anwar. Essential oils used in aromatherapy: A systemic review. *Asian Pacific Journal of Tropical Biomedicine.* 2015;5(8):601-611.
55. Eccles R. Menthol and related cooling compounds. *J. Pharm. Pharmacol.* 1994;46:618-630.
56. Peppermint Oil- Herbal summary for public, European Medicines Agency, Available:https://www.ema.europa.eu/en/documents/herbal-summary/peppermint-oil-summary-public_en.pdf. Accessed on 20 Jan 2021.
57. Keifer D, Ulbricht C, Abrams T, Basch E, Giese N, Giles M, DeFranco Kirkwood C, Miranda M, Woods J. Peppermint (*Mentha piperita*): An evidence-based systematic review by the Natural Standard Research Collaboration, *Journal of Herbal Pharmacotherapy.* 2007;7(2):91-143.
58. Haslam E, Cai Y. Plant polyphenols (vegetable tannins): Gallic acid metabolism. *Natural Product Reports.* 1994;11:41-66.
59. Andrew Pengelly. The constituents of medicinal plants. Allen & Unwin Epz Titles. 2004;2nd Ed, 116-223.
60. Anand A. Zanwar, Subhash L. Bodhankar. polyphenols in the prevention and treatment of vascular and cardiac disease, and cancer. *Polyphenols in Human Health and Disease.* 2014;2: 989-992.
61. Vatter DA, Shetty K. Biological function of ellagic acid: A review. *Journal of Food Biochemistry.* 2005; 29(3):234-266.
62. Infante R, Contador L, Rubio P, Aros D, Peña Neira Á. Postharvest sensory and phenolic characterization of Elegant Lady and Carson peaches, *Chilean Journal of Agricultural Research.* September 2011;71(3):445-451.
63. Usta C, Özdemir S, Schiariti M, Puddu PE. The pharmacological use of ellagic acid-rich pomegranate fruit, *International Journal of Food Sciences and Nutrition.* 2013 Nov;64(7):907-13.
64. Sonja Salinger-Martinovic, Vladan Cosic, Nenad Stojiljkovic, Sonja Ilic, Nikola Stojanovic, and Tijana Dencic, Impact of ellagic acid application on doxorubicin-induced cardiovascular toxicity model. *Canadian Journal of Physiology and Pharmacology.* 99(2):185-191.
65. Zhang HM, Zhao L, Li H, Xu H, Chen WW, Tao L. Research progress on the anticarcinogenic actions and mechanisms of ellagic acid. *Cancer Biol Med.* 2014;11(2):92-100.
66. Eleonora Turrini, Lorenzo Ferruzzi, Carmela Fimognari. Potential effects of pomegranate polyphenols in cancer prevention and therapy. *Oxidative Medicine and Cellular Longevity.* 2015; Article ID 938475:19.
67. Zhao Zhaohui, Moghadasian Mohammed H. Chemistry, natural sources, dietary intake and pharmacokinetic properties of ferulic acid: A review. *Food Chemistry.* August 2008;109(4):691-702.
68. Kumar Naresh, Pruthi Vikas. Potential applications of ferulic acid from natural sources: Review. *Biotechnology Reports.* December 2014; 4:86-93.
69. Gelinas Pierre, McKinnon Carole M. Effect of wheat variety, farming site, and bread-baking on total phenolics, *International Journal of Food Science and Technology.* 2006;4:329.
70. Beejmohun Vickram, Fliniaux Ophélie. Microwave-assisted extraction of the main phenolic compounds in faxseed. *Phytochemical Analysis.* July-Aug 2007; 18(4):275-282.
71. Quinde-Axtell Zory, Baik Byung-Kee. Phenolic compounds of barley grain and their implication in food product discoloration. *J. Agric. Food Chem.* December 2006;54(26):9978-9984.
72. Ludovic Rondini, Marie-Noëlle Peyrat-Maillard, Agnès Marsset-Baglieri, Claudette Berset. Sulfated ferulic acid is the main in vivo metabolite found after short-term ingestion of free ferulic acid in

- rats. J. Agric. Food Chem. 2002; 50(10):3037–3041.
73. Raoof AA, Ramtoola Z, McKenna B, Yu RZ, Hardee G, Geary RS. Effect of sodium caprate on the intestinal absorption of two modified antisense oligonucleotides in pigs. Eur J Pharm Sci. 2002 Nov;17(3):131-8.
74. Francis A. Oladimeji, Adebajo J. Adegbola, Cyprian O. Onyeji. Appraisal of bioenhancers in improving oral bioavailability: Applications to herbal medicinal products. Journal of Pharmaceutical Research International. 2018;24(4):1-23.

© 2021 Piniseti and Arora; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/77570>