



Cynodon dactylon Pers.: A Review on Forage Crop and Medicinal Herb

Papia Dutta^a and Jyotchna Gogoi^{b*}

^a Department of Chemistry, Faculty of Science, Assam Down Town University, Panikhaiti, Guwahati, Assam, 781026, India.

^b Department of Biochemistry, Faculty of Science, Assam Down Town University, Panikhaiti, Guwahati, Assam, 781026, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i58B34210

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/79165>

Mini-review Article

Received 10 October 2021
Accepted 12 December 2021
Published 15 December 2021

ABSTRACT

Cynodon dactylon is a forage crop besides being a medicinal herb which has been holding a sacred position in our Indian culture. Its survival capacity in any ecological succession has forced it to exist as medicinal plant containing a broad spectrum of secondary metabolites like Cyanidin, Luteolin, apigenin having several therapeutic uses. The review paper has been designed in coordination with research articles to compile the various properties of the plant like its thriving capacity in tropical and subtropical regions where it can alter ecosystem by effecting nutrient cycles and community composition, diverse chemical composition, and therapeutic uses to name some such as diabetes, atherosclerosis etc., which could be used as an area of further research for mankind and environmental benefits.

Keywords: *Cynodon dactylon*; forage crop; medicinal herb; cyanidin; luteolin.

1. INTRODUCTION

C. dactylon is a C₄ grass currently included in invasive plant list of Global Compedium of weeds

[1], widely distributed in tropical and subtropical regions of the world. It is a sod-forming, fast-growing grass that spreads by seeds, multiple stolons and rhizomes and colonizes

*Corresponding author: E-mail: gogoijyotchna@gmail.com;

rapidly new areas forming dense mats. It has strong potential to alter ecosystem functioning by altering hydrological cycles, fire regimes, nutrient cycles, biophysical dynamics, and community compositions [2]. Despite being categorised as an invasive plant; it has several medicinal applications and has attracted a huge amount of research especially in the field of pharmacology.

C. dactylon is one of the auspicious herbs of Ayurveda which constitutes the group 'Dasapushpam' [3]. It is a sacred plant in Indian culture next to *Ocimum* [4], mention of which has also been incorporated in early Graeco-Roman pharmacopias due to its support to cattle rearing [5]. It belongs to Poaceae family, having tinged purple colour stems, and slightly flattened in shape (Fig. 1).

2. COMMON NAME AND SYNONYMS of *C. dactylon*

C. dactylon has other two scientific names: *Panicum dactylon* and *Capriola dactylon*. It is also known by various names across the globe: in India it is called as Durva, Dhoob, Dubo, Indian doab and other English names are Bermuda grass, Bahama grass, Couch grass, Dog's tooth grass, Devil's grass, Arugampul, Scutch grass, and Grama. It is not a native grass of Bermuda but there it is found in abundance and known as crab grass [3,6].

3. *C. dactylon*: POTENTIAL FORAGE CROP

C. dactylon is distributed throughout the southernmost portions of the United States and region having similar climates worldwide. It cannot withstand freezing temperatures so, its propagation in colder belt is limited. It is

abundant in some continental climates, such as the mid-South, Texas, Oklahoma, where cold winters exist. Its growth begins only in the late spring and is maximized under the hot conditions of summer. Its growth could be established by using seed or stolon cuttings. This plant can tolerate continuous grazing therefore, over sown with other cool-season annual grasses (like ryegrass) to improve pasture production in the fall, winter and early spring periods [7].

C. dactylon has good growth rate in sandy loam soils but it can also thrive in wide range of soil types from sand to clay. On lighter soil condition, it propagates by rhizomes and stolons. It has increased establishment rate on clay soils because rhizomes and stolons cannot spread freely. The rate may however get decreased due to weed invasion but once, *C. dactylon* establishes strongly on clay soil, then it starts growing profusely.

C. dactylon is considered a forage crop that is tolerant of close grazing and need minimum management. It also acts as alternative for tall fescue during summer months. It is a versatile grass which can be grazed, hayed, and grown together with cool-season plants during dormant season. When season becomes favourable for its growth then, it becomes difficult to grow this plant with other companion forages. In some areas, alfalfa is grown with this grass. Also, Clovers could be grown with this plant, but elevated management is required for the establishment of legume and their persistence. To use this plant as forage crop in all grazing systems, one must determine the proper stocking rate of this plant under grazing management. This plant has great potential in managing grazing animals, forage production would be better if rotational grazing is preferred.




Fig. 1. *Cynodon dactylon*

4. SACRED BELIEF

In Assam also, it holds a sacred position with medicinal qualities. It is commonly called Dubori bon in Assam, many poems and novel has mention of this name. In Hindu rituals like puja and yagya, this grass is used as puja offerings to the Gods. The priest makes a ring with this grass and wear on his finger throughout the religious rituals. It is believed that the grass has purifying effect. In Ganesha temple, this grass is used as offering. As per Hindu belief, Durva grass was created by churning the mountain Mandara for Amrita (attain immortality). This churning of mountain took place between Devatas and Asuras. There is also another belief that this grass grew from the drops of Amrita which fell when Devatas were drinking it.

5. TRADITIONAL MEDICINE

C. dactylon is a weed cum medicinal plant. According to Ayurvedic Pharmacopoeia, it is pungent and bitter in taste which bears a characteristic fragrance and has cold potency. According to Unani system of medicine, this plant possesses a sharp, hot taste and pleasant odour. The aerial parts and rhizomes of the plant have been reported to be used in cardio protection, besides possessing antibacterial, antimicrobial, antioxidant, wound healing, anti-diabetic, and diuretic effects. Also, it is traditionally used for jaundice, as astringent, to stop bleeding in piles and skin infections in some parts of India namely, West Bengal, Assam, Manipur, and Mizoram [7]. Furthermore, traditional healers use it for purifying the blood, anuria, biliousness, conjunctivitis, diarrhoea, gonorrhoea, itches, Leucorrhoea, Irregular menstruation, and stomach-ache [8]. In South India, its juice is commonly consumed as health drink early in the morning to restore a healthy life. In Ayurvedic medicine, the juice of the plant could be used for treatment of hysteria, epilepsy, and insanity [9].

6. CHEMICAL CONSTITUENTS OF *C. dactylon*

Whatever might be the Hindu belief for sacred plant, the presence of phytochemicals in *C. dactylon* defines the pharmacological role of the plant. GC-MS analysis of ethanol extract of plant showed presence of glycerin; stigmasterol; gamma sterol; 1-Hydrazino-2-propanol; n-Hexadecanoic acid; Hexadecanoic acid ethyl ester; 9,12,15-Octadecatrienoic acid (Z, Z, Z); 1-

Triacantanol, 2-Dodecen-1-yl (-) succinic anhydride; 3,7,11,15-tetramethyl-2-hexaden-1-ol (phytol) [10].

Preclinical studies of *C. dactylon* have shown that the oral administration at a dose of 20 mg/kg/body wt decreased the adjuvant-induced inflammatory response. It also reduced oxidative stress and ameliorated the arthritic changes to normal conditions. When compared with the control, extract of *C. dactylon* decreased the levels of inflammatory mediators, myeloperoxidase, nitrite, C-reactive protein, ceruloplasmin, and reactive oxygen species. Simultaneously, increased level of antioxidant system like catalase, superoxide dismutase, glutathione peroxidase, glutathione, vitamins C and E has been reported [3].

The plant is also used in treating dropsy [11], syphilis [12], and diabetes [13]. Studies have shown that active principles of *C. dactylon* found in ethanolic extracts exhibited antioxidant activity, namely, Cyanidin, beta-carotene, and tritacin [14].

Studies have shown that protein fraction of *C. dactylon* could sensitized the humoral and cell mediated immunity [15]. Research work conducted in year 1999 [16], revealed that the administration of *C. dactylon* could significantly increase the antibody responsiveness in mice to sheep RBCs by increasing the counts of macrophages and antibody-synthesizing cells. Another study conducted in 2009, also proved that this plant could increase humoral antibody response against antigen. Also, it could inhibit the release of TNF- α , IL-6, and IL-1 [17]. A study reported its anti-inflammatory action by observing increased IL 10 secretion which in turn suppressed proinflammatory cytokines [18]. This plant not only could modulate immunity in mice, but also when *Catla catla* fish was supplied with food containing *C. dactylon* extract, a nonspecific immunity could be observed which protected it from bacterial infection caused by *Aeromonas hydrophila* [19].

This plant contains a minor storage protein tritacin which is reported with antioxidant activity. Tritacin is a storage protein of wheat endosperm which is nutritionally rich due to the presence of unique lysine- rich decapeptide repeat motif inserted in the hyper variable region of this gene [20]. It belongs to legumin super family (11-12S globulins) storage protein. It is predominant in seeds of legumes and some cereals such as rice and oat [21].

It has been also reported that this plant contains very important metabolite called cyanidin (Fig 2). It is a plant pigment of anthocyanidin origin and very potent chemopreventive agent. Most of berries display the colours due to Anthocyanins, particularly glucosides and galactosides of cyanidin, peonidin, delphinidin, petunidin and malvidin. During ripening season, pigment secretion increases whereas decreased secretion has been seen for polyphenols (-)-epicatechin, (+)-catechin and dimeric proanthocyanins [22].




Fig. 2. Cyanidin

C. dactylon exhibits antioxidant properties due to the presence of phytochemicals like beta carotene which has been reported in this plant [14]. Beta-carotene belongs to the group of carotenoids consisting of isoprene units. The presence of conjugated pi-system in the long carbon chain of beta-carotene acts as a chromophore responsible for its specific color [23]. It is the most abundant form of carotenoid, and it is a precursor of vitamin A. Beta-carotene is composed of two retinyl groups. It can be found in yellow, orange, and green leafy vegetables and fruits [24].

β -Carotene (Fig. 3) is an antioxidant that has greater efficacy against the reactive oxygen species, i.e., singlet oxygen (1O_2) compared to vitamin E and vitamin C [25]. Feeding β -carotene in the diet of C57BL/6J mice beginning at 1 month or at 20 months of age for the duration of the life span increased the serum concentration of β -carotene by 60% but did not change the β -carotene content of heart, liver, or kidney and neither significantly affected the mean nor the maximal life span [26].

Many research works have highlighted different pharmacological properties such as antioxidant, antiarthritic, anti-inflammatory, antimicrobial, antidiuretic, and anti-atherosclerotic property. In

2014, antiarthritic study revealed that *C. dactylon* could effectively improve WBCs, ESR, C-reactive protein (CRP) and reduce TNF alpha level. Also, the study mentioned that the plant could protect arthritic joints by improving bone lesions [31]. Another study conducted in 2009 showed that when *C. dactylon* dose of 20mg/kg/body weight were given orally to arthritic rats after adjuvant injection as compared to dose treatment of 3mg/kg/body weight, a significant improvement in the inflammatory response, oxidative stress and ameliorated arthritic changes to reach normal state was observed [3]. Early atherosclerotic changes in the vessel wall would be prevented by treating the high cholesterol diet rats with ethanolic extract of *C. dactylon* [32]. Also, reports have shown that this plant could show a strong protective effect on right heart failure by giving positive inotropic action and improving cardiac functions [33]. A study revealed that aqueous decoction of *C. dactylon* could reduce stone formation at dose of 200mg/kg/ body weight as compared to 12.5 and 50mg/kg/body weight [34]. Also, this plant has the capacity to prevent and eliminate Calcium oxalate deposition into kidneys of rat which supports the traditional belief of the plant to be used in treatment of kidney stones [35]. Even fractional analysis of this grass has supported the traditional belief of kidney stone treatment [36]. In 2015, 2017 and 2018, different studies were conducted which promoted different stages of wound healing like angiogenesis [37]; collagenesis [38] and Collagen Silica Cynodon extract scaffold for treating wounds [39].




Fig. 3. β - carotene

8. CONCLUSION

C. dactylon, a sacred plant of Indian culture and globally used as forage crop. The review could conclude that many research articles have shown beneficial roles of the plant to the ecosystem and medicine. Study has highlighted reason for use of this plant as forage crop. As per literature on traditional belief, *C. dactylon* is very good herb for gynaecological issues like Leucorrhoea, irregular menstruation. However,

no report has been found in this regard which can validate its use in gynaecology treatments.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Randall RP. A global compendium of weeds. Dpt. Agric. Food Western Australia 2012;1124.
2. Burns JC, Fisher DS. "Coastal" and "Tifton 44" Bermudagrass Availability on Animal and Pasture Productivity. *Agronomy J.* 2008;100(5):1280-88.
3. Sindhu G, Ratheesh M, Shyni GL, Helen A. Inhibitory effects of *Cynodon dactylon* L. on inflammation and oxidative stress in adjuvant treated rats. *Immunopharmacol Immunotoxicol.* 2009;31(4):647-53.
4. DURVA (Dub) (*Cynodon dactylon*) ; 2011. Available: www.herbalcureindia.com/herbs/durva.htm
5. Beck P, Jennings J, Rogers J. Management of pastures in the upper south: The I-30 and I-40 corridors. In: Management strategies for Sustainable Cattle production in Southern Pastures. Academic Press. 2020;189-226.
6. Pal D, Pandab K. Evaluation of Anthelmintic Activities of Aerial parts of *Cynodon dactylon* Pers. *Ancient Sci Life.* 2010;30(1): 12-3.
7. Pitman WD. Forage systems for warm humid areas. In: Forages- The Science of Grassland Agriculture. Sixth ed. Blackwell Publishing. 2007;2:303-12.
8. Bharati D, Sonawane SA, Amrutkar MP, Undale VR, Wankhade AM, Bhosale AV. *In Vivo* Evaluation of Antidiarrhoeal activity of aerial parts of *Cynodon dactylon* Linn. Pers. (Poaceae). *Pharmacol Online.* 2009; 2:1-11.
9. Chellaiah M, Marriappan A, Nagappan R, Savarimuthu I. Medicinal plants used by traditional healers in Kancheepuram District of Tamil Nadu. *Indian J. Ethnobiol. Ethnomed.* 2006;2(43):1-10.
10. Rawal JR, Sonawani PR. Determination of Bioactive Components of *Cynodon dactylon* by GC-MS Analysis & it's *In Vitro* Antimicrobial Activity. *Int J Pharm Life Sci.* 2016;7(1):4880-5.
11. Kesari AN, Gupta RK, Singh SK, Diwakar S, Watal G. Hypoglycemic and antihyperglycemic activity of *Aegle marmelos* seed extract in normal and diabetic rats. *J Ethnopharmacol.* 2006; 107:374-9.
12. Provin TL, Wright AL, Hons FM, Zuberer DA, White RH. Seasonal dynamics of soil micronutrients in compost-amended bermudagrass turf. *Bioresources Technol.* 2008; 99(7):2672-9.
13. Singh SK, Rai PK, Jaiswal D, Rai DK, Sharma B, Watal G. Protective effect of *Cyanodon dactylon* against STZ induced hepatic injury in rats. *J Ecophysiol Occup Health.* 2008;8(3&4):195-9.
14. Auddy B, Ferreira M, Blasina F, Lafon L, Arredondo F, Dajas F, Tripathi PC, Seal T, Mukherjee. Screening of antioxidant activity of three Indian medicinal plants, traditionally used for the management of neurodegenerative diseases. *J. Ethnopharmacol.* 2003;84(2-3):131-8.
15. Santhi R, Annapoorani S. Efficacy of *Cynodon dactylon* for immunodulatory activity. *Drug Invention Today.* 2010; 2(2):112-4.
16. Shah VK, Zalavadiya VI, Sheth NR, Raval MK, Dudhrejiya Av, Golwala DK. Standardization and Assessment of Preformation Parameters of Rasayana Tablet. *Int. J. Pharma. Sci Drug Res.* 2010;2(1):58-62.
17. Mangathayaru K, Umadevi M, Reddy CV. Evaluation of the immunomodulatory and DNA protective activities of the shoots of *Cynodon dactylon*. *J Ethnopharmacol.* 2009;123(1):181-4.
18. Moore KW, de Waal Malefyt R, Coffiman RLO, Garra A. Interleukin -10, and the Interleukin-10 receptor. *Annu Rev Immunol.* 2001;19(1):632-3.
19. Kaleeswaran B, Ilavenil S, Ravikumar S. Dietary supplementation with *Cynodon dactylon* (L) enhances innate immunity and disease resistance of Indian major carp,

- Catla catla* (Ham). Fish Shellfish Immunol. 2011;31(6):953-62.
20. Arcalis E, Marcel S, Altmann F, Kolarich D, Drakakaki G, Fischer R, Christou P, Stoger E. Unexpected deposition patterns of recombinant proteins in post-endoplasmic reticulum compartments of wheat endosperm PlantPhysiol. 2004;136(3):345-7-3466.
 21. Yadav D, Singh NK. Wheat Triticin: A Potential Target for Nutritional Quality Improvement. Asian J Biotectnol. 2011; 3(1):1-21.
 22. Evans WC, Evans D. Phenols, and phenolic glycosides. In: Trease and Evans' Pharmacognosy. W.B. Saunders 2009;219-62.
 23. Flora G, Mittal M, Flora SJS. Medical Countermeasures-Chelation Therapy. In: Handbook of Arsenic Toxicology. Academic Press. 2015;589-626.
 24. Pattani KM. Chemoprevention of Head and Neck Cancer. In: Early diagnosis and treatment of cancer series: Head and Neck Cancer Emerging Perspectives. Academic Press. 2010;81-93.
 25. Roberts JE, Dennison J. The Photobiology of Lutein and Zeaxanthin in the Eye. J Ophthalmol; 2015.
Available:<https://doi.org/10.1155/2015/687173>
 26. Kaur S, Sharma R, Sarangal V, Kaur N, Prashar P. Evaluation of anti-inflammatory effects of systemically administered curcumin, lycopene and piperine as an adjunct to scaling and root planing: A clinical study. AYU. 2017;38(2):117-21.
 27. Sumathi T, Chandrasekar S, Balasubramanian RK, Devarajulu NN. Protective role of *Cynodon dactylon* in ameliorating the aluminium-induced neurotoxicity in rat brain regions. Biol Trace Elem Res. 2011;144(1-3):843-53.
 28. Pattanashetti LA, Patil BMP, Hegde HV, Kangle RP. Potential ameliorative effect of *Cynodon dactylon* (L.) pers on scopolamine-induced amnesia in rats: Restoration of cholinergic and antioxidant pathways. Indian J Pharmacol. 2021;53(1): 50-9.
 29. Jarald EE, Joshi SB, Jain DC. Antidiabetic activity of aqueous extract and non-polysaccharide fraction of *Cynodon dactylon* Pers. Indian J Exp Biol. 2008; 46(9):660-7.
 30. Annapurna HV, Babu A, Natesan R, Kallur PA, Pemaiah B, Sethuraman S, Mahadevan V, Arumugam NJ. Isolation and *in silico* evaluation of antidiabetic molecules of *Cynodon dactylon* (L.). J Mol Graphics Modelling. 2013;39:87-97.
 31. Bhangale J, Acharya S. Antiarthritic activity of *Cynodon dactylon* (L.) Pers. Indian J Exp Biol. 2014;52(3):215-22.
 32. Pashaie B, Hobbenaghi R, Malekinejad H. Anti-atherosclerotic effect of *Cynodon dactylon* extract on experimentally induced hypercholesterolemia in rats. Vet Res Forum Summer. 2017; 8(3): 185-93.
 33. Garjani A, Afroozian A, Nazemiyeh H, Najafi M, Kharazmkia A, Maleki-Dizaji N. Protective effects of hydroalcoholic extract from rhizomes of *Cynodon dactylon* (L.) Pers. on compensated right heart failure in rats. BMC Complement Altern Med; 2009.
DOI: 10.1186/1472-6882-9-28.
 34. Golshan A, Hayatdayoudi P, Hadjzadeh MAR, Rad AK, Roshan NM, Abbasnezhad A, Mousavi SM, Pakdel R, Zarei B, Aghaee A. Kidney stone formation and antioxidant effects of *Cynodon dactylon* decoction in male Wistar rats. Avicenna J Phytomed. 2017;7(2):180-90.
 35. Atmani F, Sadki C, Aziz M, Mimouni M, Hacht B. *Cynodon dactylon* extract as a preventive and curative agent in experimentally induced nephrolithiasis. Urol. Res. 2009;37(2):75-82.
 36. Khajavi Rad A, Hadjzadeh MA, Rajaei Z, Mohammadian N, Valiollahi S, Sonei M. The beneficial effect of *Cynodon dactylon* fractions on ethylene glycol-induced kidney calculi in rats. Urol J 2011;8(3):179-84.
 37. Soraya H, Moloudizargari M, Aghajanshakeri S, Javaherypour S, Mokarizadeh A, Hamedeyazdan S, Ghaleh HEG, Mikaili P, Garjani A. Angiogenic effect of the aqueous extract of *Cynodon dactylon* on human umbilical vein endothelial cells and granulation tissue in rat. Daru. 2015; 23(1): 10.
DOI: 10.1186/s40199-015-0093-x.
 38. Biswas TK, Pandit S, Chakrabarti S, Banerjee S, Poyra N, Seal T. Evaluation of *Cynodon dactylon* for wound healing activity. J Ethnopharmacol. 2017;197:128-37.
 39. Perumal RK, Gopinath A, Thangam R, Perumal S, Masilamani D, Ramadass SK, Madhan B. Collagen-silica bio-composite

enriched with *Cynodon dactylon* extract for
tissue repair and regeneration. Mater Sci

Eng C Mater Biol Appl. 2018;92:297-306.

© 2021 Dutta and Gogoi; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/79165>