



Effect of Provenance Variations on the Growth and Development of *Terminalia ivorensis* (A Chev)

M. O. Ojo¹ and Ajayi Ezekiel^{1*}

¹Department of Forestry and Wildlife Management, Adekunle Ajasin University, Akungba Akoko, Ondo State, Nigeria.

Authors' contributions

This work was carried out in collaboration between both authors. Author AE designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors MOO managed the analyses of the study and managed the literature searches. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CJAST/2019/v37i130274

Editor(s):

(1) Dr. Aydin Unay, Professor, Department of Field Crops, Faculty of Agriculture, University of Aydin Adnan Menderes, Turkey.

Reviewers:

(1) Adelalu Tope Gabriel, Taraba State University, Nigeria.

(2) Vinay Kumar Pandey, India.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50867>

Original Research Article

Received 30 May 2019
Accepted 14 August 2019
Published 21 August 2019

ABSTRACT

Indigenous tree species like *Terminalia ivorensis* has multiple uses; from timber products to medicinal condiments for treating different ailments. This species is self-incompatible. Thus, habitat protection is a key to its survival and sustenance. This study evaluated the effect of provenance variations on the growth and development of *T. ivorensis*. Three provenances of *T. ivorensis* were selected; FRIN Arboretum, Onigambari Forest Reserve and J1 Forest Reserve. Thirty (30) seedlings of *T. ivorensis* were selected and marked at each provenance for assessment of growth parameters for a period of twelve weeks. The growth parameters included: seedling height, collar diameter, leaf count and leaf area. Also, soil samples were collected from each provenance to test for nutrient composition. Results showed that J1 Forest Reserve had best performance across the growth parameters assessed with 10.4 cm plant height, 2.02 mm collar diameter, 11 leaves count and 20.5 cm² leaf area respectively. This growth parameter result differs significantly from the other two provenances. This was also attributed to the nutrient offered by J1 Forest Reserve with 1.18% and 2.20% of organic carbon and organic matter in the soil respectively. Also, the available phosphorus which is involved in photosynthesis, respiration, energy storage and transfer, cell division and enlargement as well as promotion of root formation was about 15.7 mg/kg in the soil

*Corresponding author: E-mail: ezekielajayi33@gmail.com, ezekiel.ajayi@aau.edu.ng;

and is considered high because it is higher than the critical level of 8.50mg/kg. J1 Forest Reserve is recommended as the best provenance for raising *T. ivorensis* amongst assessed natural habitats.

Keywords: Collar; *ivorensis*; leave; parameter; provenance; *Terminalia*.

1. INTRODUCTION

Terminalia ivorensis belongs to the family combretaceae and it is an important timber species [1] recognized in Nigeria with its various uses. *Terminalia ivorensis* also plays a major role in increasing soil fertility where fall off leaves are decomposed and add more nutrients to the soil [2]. This species is widely used for medicinal purposes in Africa [3,4]. This tree species if protected from logger can attain heights of up to 50 m and girth of 500 cm. It has been biologically stated that *T. ivorensis* is self-incompatible; the flowers are bisexual, its selection and breeding started in the 1960s in Africa. [5] Stated that the trees with superior growth rate and stem form have been selected and clone, banks have been established in this species. Though, the interval between the opening of the leaf buds and flowering usually takes 3-4 weeks where the flowers are fertilized by insects. Fruiting, which begins in December, is abundant from January to March [5].

The trees sapling usually requires 4 – 5 years before attaining 1-1.5 m height, compared to other indigenous timber species with fast growth. For example, even *T. superba* from the same genus could attain 1.8 m in height within one year [6]. *Terminalia ivorensis* is used for construction and other timber-based uses [7]. It's also can be used for reforestation and afforestation species especially where wood-base timber are needed as well as for forest ecosystem replenishment [8]. However, this species occurs naturally in evergreen and moist semi-deciduous forests ecosystem, where large trees are most common in low density localities or lowlands [9]. It is most abundant in the transition zone between humid semi-deciduous and evergreen forests. It is found in rainforest ecosystem but predominantly it's a tree of seasonal forest zones [5].

As one of the principal timber species of West Africa countries, *T. ivorensis* is widely harvested from natural forest and has been introduced into many other tropical countries as a promising timber plantation species. It is also grown as a shade tree in cocoa plantations in Nigeria [5]. It

has been reported that the species is threatened by habitat loss and poor regeneration, and attempts at plantation growth have generally failed through frequent diebacks. The species has also been classified as 'vulnerable' by the IUCN [5]. Despite these facts, plantation of *T. ivorensis* is still scarce in the country. With so much emphasis today on ecosystem management and maintenance of natural forests, sustainable artificial regeneration of tree species for large-scale plantation development has become expedient to meet up with wood-based need in the country as well as environmental sustenance in a world of climate change effects where oil exploration has led to serious deforestation and degradation of several valuable forest species. However, not much work has been done on *T. ivorensis* in Nigeria. Although, some indigenous tree species like *P. biglobosa* and *T. superba* etc. do not experience much difficulty during germination, but *T. ivorensis* germinate with great difficulty [5].

There is however a dearth of information on the best conditions for the propagation of the species. With deliberate efforts being made by the Nigeria Government to conserve some valuable indigenous tree species both *in-situ* and *ex-situ*, efforts have been made without much success recorded. If germination of this tree species encountered difficulty then alternative knowledge for it would go a long way in encouraging its plantation establishment.

Consequently, it is pertinent to ascertain the best provenance of growing *T. ivorensis* to ensure better performance of the species. So this study is aimed at assessing the effect of provenance variation on the growth and development of *Terminalia ivorensis*.

1.1 Study Area

This study was carried out in three *T. ivorensis* provenances which are: Onigambari Forest Reserve which lies approximately between latitude 7°7'60" N and longitude 3°49'60" E; Forestry Research Institute of Nigeria (FRIN) Arboretum lies approximately between latitude 7°23'28.68" N and longitude 3°51'46.08" E; and

J1 Forest Reserve lies approximately between latitude 7°0'0" N and longitude 4°15'0" E respectively. The climatic condition of these three provenances is dominated by rainfall ranging from 1200 mm to 1500 mm annually and average temperature 35°C, and the average relative humidity ranges between 80 – 85% [10].

2. METHODS

Ninety uniform seedlings of *T. ivorensis* were identified and marked for measurement of growth performance in a period of 12 weeks (3 months). The uniformity of the seedlings were determined by selecting seedlings with height 8.0 cm, collar diameter of 1.0 mm and 7 leaves count stage. Each seedling was marked and the coordinator of the location recorded with a GPS receiver. Thirty (30) seedlings were selected from each provenance (i.e. Onigambari Forest Reserve, FRIN Arboretum and J1 Forest Reserve). Also, soil sample was collected with soil auger at each provenance to determine nutrient composition of the three provenances. Growth parameters (seedling height, collar diameter, leaf count and leaf area) were measured for three months. Seedling height was measured using graduated metric ruler; collar diameter was measured with vernier calliper, leaf count was done visually and leaf area was determined using formula adopted by Agboola and Ayodele [11].

2.1 Experimental Design and Data Analysis

Randomise Complete Block Design (RCBD) was adopted as experimental design since there are two sources of variations. Variation due to provenance as treatment and growth parameter as block since the objective of the study is to assess effect of provenance variation on the growth and development of *T. ivorensis*. The data collected were analysed with Analysis of Variance (ANOVA)

3. RESULTS AND DISCUSSION

The results from the ninety seedlings selected for this study were collated and the mean from each parameter were reported see Table 1. J1 Forest Reserve had the highest plant height with 10.4 cm (0.104 m), followed by Onigambari Forest Reserve with 9.11 cm (0.901 m) and FRIN Arboretum had the least plant height with 8.45 cm (0.805 m). Also, the leaf area; 20.5 cm² was offered by J1 Forest Reserve, Onigambari Forest Reserve followed closely with 20.2 cm² and 19.2

cm² was recorded for FRIN Arboretum (see Table 1) for details on collar diameters and leaf count results.

There were significant differences among the different provenances at $P > 0.05$ (i.e. Onigambari Forest Reserve, FRIN Arboretum and J1 Forest Reserve) as well as the growth parameters (plant height, collar diameter, leaf count and leaf area) as adopted for this study see Table 2 for details.

The mean of the different provenances (treatments) were separated using Least Significant Difference (LSD). The result showed that there were significant differences between the means ($P > 0.05$). J1 Forest Reserve mean differed significantly from both FRIN Arboretum and Onigambari Forest Reserve. Although, FRIN Arboretum and Onigambari Forest Reserve means were not significantly different from each other ($P > 0.05$) (see Table 3) for details.

3.1 Soil Analysis

Soil sample obtained from different provenances were subjected to laboratory analysis. The result shows variation across the nutrients composition of the provenance which resulted in the variations in the growth parameters.

3.2 Onigambari Forest Reserve

Table 4 below showed the laboratory analysis of soil sample collected from Onigambari Forest Reserve. The soil analysis showed that the textural class of the soil is sandy-loam. Both the organic carbon and organic matter in the soil were low with 0.88% and 1.51% respectively; this is because organic matter was above the critical level of 2.00 as reported by [12]. Nitrogen which helps in the formation of amino-acids and the building of protein is necessary for plant cell division. However, based on the result, total nitrogen available in the soil is 0.07% which is below the critical level of 0.15% [12]. Also, the inadequacy of nitrogen had an effect on the growth of Onigambari Forest Reserve seedlings by causing a decrease in their growth and also chlorosis which could be seen in the collar diameter of the seedlings. The available phosphorus which is involved in photosynthesis, respiration, energy storage and transfer, cell division and enlargement is about 2.16% in the soil which is considered low because it is below the critical level of 8.50% [12].

Table 1. Results of growth parameter grouped as treatments and blocks

Treatment/ Block	Onigambari forest reserve	FRIN arboretum	J1 forest reserve
I. Plant height (cm)	9.11	8.45	10.4
II. Collar diameter (mm)	1.14	1.93	2.02
III. Leaf count (visual)	10	9	11
IV. Leaf Area (cm ²)	20.2	19.2	20.5

Table 2. Tests of between-subjects effects (ANOVA table)

Dependent variable: data					
Source	Type III sum of squares	Df	Mean square	F	Sig.
Corrected Model	509.172 ^a	5	101.834	361.154	.000
Intercept	1259.725	1	1259.725	4467.595	.000
Provenance	3.671	2	1.836	6.510	.031
G. parameter	505.501	3	168.500	597.584	.000
Error	1.692	6	.282		
Total	1770.590	12			
Corrected Total	510.864	11			

a. R Squared = .997 (Adjusted R Squared = .994)

Table 3. Least significant differences (Multiple comparisons)

(I) Treatment	(J) Treatment	Mean difference (I-J)	Std. error	Sig.	95% confidence interval	
					Lower bound	Upper bound
FRIN Arbo	JI FR	-1.3350 [*]	.37548	.012	-2.2538	-.4162
	Onigambari	-.4675	.37548	.260	-1.3863	.4513
JI FR	FRIN Arbo	1.3350 [*]	.37548	.012	.4162	2.2538
	Onigambari	.8675	.37548	.060	-.0513	1.7863
Onigambari	FRIN Arbo	.4675	.37548	.260	-.4513	1.3863
	JI FR	-.8675	.37548	.060	-1.7863	.0513

Based on observed means. The error term is Mean Square (Error) = .282

*. The mean difference is significant at the 0.05 level

Table 4. Nutrients compositions of onigambari forest reserve

S/N	Soil properties	Value
1	Total organic carbon	0.88%
2	Total organic matter	1.51%
3	Total Nitrogen	0.07%
4	Phosphorus	2.16 mg/kg
5	Magnesium	3.33 mol/kg
6	Zinc	38.2 mg/kg
7	Iron	14 mg/kg

Though, micronutrients are high with Iron (Fe) having 14 mg/kg, which is above the critical level of 3.50 mg/kg, which is important in the

production of chlorophyll, and also component of many enzymes associated with the energy transfer and fixation of lignin formation [12]. Zinc

Table 5. Nutrients compositions of J1 forest reserve

S/N	Soil properties	Value
1	Total organic carbon	1.18%
2	Total organic matter	2.20%
3	Total Nitrogen	0.06%
4	Phosphorus	15.7 mg/kg
5	Magnesium	7.6 mol/kg
6	Nitrogen	0.76 mol/kg
7	Zinc	17 mg/kg
8	Iron	73.5 mg/kg

Table 6. Nutrients compositions of FRIN arboretum

S/N	Soil properties	Value
1	Total organic carbon	0.38%
2	Total organic matter	0.65%
3	Total Nitrogen	0.03%
4	Phosphorus	7.896 mg/kg
5	Magnesium	13.35 mol/kg
6	Zinc	19.8 mg/kg
7	Iron	71.6 mg/kg

(Zn) is essential for protein synthesis and growth regulation had a positive effect on the seedlings in terms of leaf production and leaf area, 38.2 mg/kg was obtained and it's above the critical level of 1.0 mg/kg [12].

3.3 J1 Forest Reserve

The Table 5 showed the laboratory analysis of soil sample collected from J1 Forest Reserve. The analysis revealed that the textural class of the soil is sandy-loam. Both the organic carbon and organic matter in the soil are high with 1.18% and 2.20% respectively; this is because organic matter is above the critical level of 2.00% [12]. Nitrogen which helps in the formation of amino-acids and the building of protein is necessary for plant cell division. However, based on the result below, total nitrogen available in the soil is 0.06% which is below the critical level of 0.15%, thereby having an effect on the growth of plants by causing stunted growth and chlorosis which was vividly seen among the seedlings of J1 forest reserve through the growth parameters assessed. The available phosphorus which is involved in photosynthesis, respiration, energy storage and transfer, cell division and enlargement, which also promotes root and formation is about 15.7 mg/kg in the soil and is considered high because it is higher than the critical level of 8.50 mg/kg [12]. In relation to the growth of the seedlings, it helped in boosting the

growth of the seedlings in the terms of stem diameters when compared to the seedlings from other provenances.

Also, Iron (Fe), which is involved in the production of chlorophyll, which is also a component of many enzymes associated with the energy transfer and fixation of lignin formation in plants is available as 3.75 mg/kg which is considered high because it is above the critical level of 3.50 mg/kg [12]. Zinc which took part in the regulation of growth among the seedlings of its provenance and essential for protein synthesis as well as growth regulation is about 17.0 mg/kg in the soil which is above the critical level of 8.50 mg/kg [12].

3.4 FRIN Arboretum

Table 6 showed the laboratory analysis of soil sample collected from FRIN Arboretum. The soil analysis revealed that the textural class of the soil is sandy-loam. Both the organic carbon and organic matter in the soil are low with 0.38% and 0.65% respectively and this organic matter is above the critical level of 2.00% [12]. Total Nitrogen which helps in the formation of amino-acids is 0.03% which is below the critical level of 0.15%. Based on the performance of the seedlings from this provenance, the inadequacy of nitrogen had a serious effect on the growth of plants in terms of leaf area, collar diameter and

plant height when compared to the seedlings of other provenances. Phosphorus (P) which is involved in photosynthesis, respiration, energy storage and transfer is high about 7.90% in the soil which is below the critical level of 8.50% [12].

Iron (Fe), which helps in the production of chlorophyll also attributed to the seedlings growth in terms of collar diameter. Copper (Cu), which is available as 18 mg/kg, is above the critical level of 3.50 mg/kg helped in providing necessary carbohydrate and nitrogen metabolism and it also on the other hand help preventing wilting plants. Also, Zinc (Zn) which is essential for protein synthesis and growth regulation had a positive effect on the seedlings by enhancing their growth in terms of leaf area and leaf production is about 71.6 mg/kg in the soil which is above the critical level of 8.50 mg/kg [12].

4. CONCLUSIONS AND RECOMMENDATIONS

There are variations in growth parameters of seedlings from the three provenances assessed for this study. This can be seen and correlated with soil sample analysis results from the provenances. The result of the soil analysis showed that J1 Forest Reserve had the highest nutrients and this resulted in the growth parameter been the best for the *Terminalia ivorensis* seedlings. There were significant differences amongst the provenances as well as growth parameters investigated.

As a result of means separation, J1 Forest Reserve differed significantly from the other two provenances and is recommended as the best provenance for raising *Terminalia ivorensis*. Therefore, for growth and development of *Terminalia ivorensis* on large scale, J1 Forest Reserve is the best amongst location where this species are found in southwestern, Nigeria.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Foli EG. *Terminalia ivorensis* A. Chev. In: Lemmens RHMJ, Louppe D and Oteng-Amoako AA. (Editors). PROTA (Plant Resources of Tropical Africa / Ressources végétales de l'Afrique tropicale), Wageningen, Netherlands; 2009.
2. Norgrove L, Hauser S. Yield of plantain grown under different tree densities and 'slash and mulch' versus 'slash and burn' management in an agrisilvicultural system in southern Cameroon. *Forest Ecology and Management*. 2002;166:261-270.
3. Coulibaly K, Zirihi GN, Guessennd-Kouadio N, Oussou KR, Dosso M. Antibacterial properties of trunk barks of *terminalia ivorensis* (Combretaceae), a commercial and medicinal specie, on some methicillin-resistant *Staphylococci* spp strains. *African Health Sciences*. 2014;14(3):753-756.
4. Masoko P, Picard J, Eloff JN. Antifungal activities of six South African *Terminalia* species (Combretaceae). *Journal of Ethnopharmacology*. 2005;75:45-50.
5. Orwa C, Mutua A, Kindt R, Jamnadass R, Simons AJ. *Agroforestry database: A tree reference and selection guide version 4.0*; 2009.
Available: https://www.researchgate.net/publication/258996453_Agroforestry_Database_A_Tree_Reference_and_Selection_Guide_version_40
6. Osei-Begyina A. Effect of inorganic (NPK) fertilizer on the growth rate of one-year old *diospyros mespiliformis*. B.Sc. thesis, Department of Silviculture and Forest Management, Faculty of Renewable Natural Resources, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana. 2007;40.
7. Mohammed AI, Ocholi A, Ejeh SP, Amartey YD. Compression creep model for Nigeria grown *Terminalia ivorensis* (Black afara) timber. *Nigerian Journal of Technology*. 2019;38(2):311-316.
8. Jones RK, Averre CW. Damping-off in flower and vegetable seedlings: Plant pathology extension. North Carolina State: University Ornamental Disease Note. No. 14; 2000.
9. Ibe AE, Adeyemi AA, Nwuda BN, Asiabaka CC, Onuoha GN. Seed germination and seedling growth of *Terminalia ivorensis* (A. Chev) in different growth media. *FUTO Journal Series (FUTOJNLS)*. 2015;1(2):2476-8456.
10. Clifton-Brown JC, Jones MB. The thermal response of leaf extension rate in genotypes of the grass *Miscanthus*: An

- important factor in determining the potential productivity of different genotypes. Journal of Experimental Botany. 1997;48:1573-1581.
11. Agboola AA, Ayodele OJ. Prospects and problems of using soil testing for adoption of fertilizer use in Ekiti-Akoko agricultural development project area. In processing's of workshop on appropriate technologies for farmers in Semi-Arid West Africa, Ohm HW, JG Nagy (Eds) Purdue University, West Lafayette. 1985;123-136.

© 2019 Ojo and Ezekiel; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/50867>