



Characterization and Correlation Analysis of Physical and Physico-Chemical Properties of Cambucá Fruits (*Plinia edulis*)

**Antônio Gustavo de Luna Souto^{1*}, Maria Helena Menezes Cordeiro¹,
João Paulo Gava Cremasco¹, Valtânia Xavier Nunes¹, Keise Barcelas Moraes¹
and Carlos Eduardo Magalhães dos Santos¹**

¹Department of Plant Science, Universidade Federal de Viçosa, Viçosa-MG, Brazil.

Authors' contributions

This work was carried out in collaboration between all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2019/v31i430077

Editor(s):

(1) Asst. Prof. Biljana Bojovic, Faculty of Science, Institute of Biology and Ecology, University of Kragujevac, Republic of Serbia.

Reviewers:

(1) Valdir Florencio da Veiga Junior, Military Institute of Engineering, Brazil.

(2) Mohd Zubair Tak, Pulwama.

Complete Peer review History: <http://www.sdiarticle3.com/review-history47242>

Original Research Article

Received 22 October 2018
Accepted 05 February 2019
Published 28 February 2019

ABSTRACT

Aims: This study was to characterize and correlate physical and physico-chemical properties of cambucá fruits (*Plinia edulis*).

Study Design: Experimental design was completely randomized with fifteen samples of five fruits each.

Place and Duration of Study: Experimental orchard of the Department of Plant Science, Federal University of Viçosa (UFV), located in the municipality of Viçosa, Zona da Mata of Minas Gerais during the month of February 2015.

Methodology: Fruits were sampled when 100% of peels presented yellow-orange color. The following characteristics were evaluated: longitudinal and transverse diameter, total mass of both pulp and seed, pulp color, soluble solids, titratable acidity, soluble solids/titratable acidity ratio, ascorbic acid and carotenoids contents. To verify the degree of correlation between two physico-chemical variables, a Pearson analysis was performed.

*Corresponding author: E-mail: gusluso@hotmail.com;

Results: Cambucá fruits showed average values of longitudinal and transverse diameter of 37.76 and 44.36 mm, respectively. Fruits' average mass were 44.12 g and the percentage of pulp was 82.15%. Both soluble solids and titratable acidity presented the respective average values: 10.53 °Brix, 1.34 mg of citric acid and 100 mL⁻¹ of pulp. Larger cambucá fruits presented higher pulp yield and lower acidity. The increase in ascorbic acid was positively correlated with the contents of soluble solids and carotenoids.

Conclusion: Fruits of cambucá have potential for commercialization, their characteristics are similar to those found in other fruit species native to the Myrtaceae family and already found in the fruit market.

Keywords: Analysis pearson; carotenoids; exotic fruits; myrtaceae; *Plinia edulis*.

1. INTRODUCTION

The species *Plinia edulis* is native from the Brazilian Atlantic forest, popularly known as “cambucazeiro” or “cambucá”, growing from the State of Rio de Janeiro to Santa Catarina, being currently in danger of extinction [1]. It is a tree species, with height varying from 5 to 10 meters and belongs to the family Myrtaceae, which houses a great diversity of fruit species, such as jaboticaba (*Myrciaria jaboticaba*), pitanga (*Eugenia uniflora* L.), uvaia (*Eugenia pyriformis* Cambess.), guariroba (*Campomanesia xantocarpa* O. Berg.), cerejeira-do-mato (*Eugenia involucrate* DC.), among others [2,3].

The cambucazeiro is well known in local communities where the species is found, due to its fleshy and juicy fruits, widely used in folk medicine against stomach problems and sore throat [1]. Some important substances for the pharmaceutical industry are found in its leaves, such as flavonoids, terpenoids and tannin [1,4], while the fruits have acceptable sensory characteristics and nutritious properties of interest [5,6].

Although commonly used in local medicine, only a few scientific works were made with this species, justifying the need of more studies that may allow greater exploitation of the cambucá fruit [2,4,7]. Studies related to fruit's characterization are of great relevance to this exploitation, given new marketing possibilities for its potentialities. Native fruits from the Atlantic forest usually have good levels of vitamins, mineral salts and antioxidant composites [8,9,10,11].

Therefore, the lack of information that may enhance the potentiality of the cambucazeiro through fruits characterization, thus allowing the exploitation and consequently the perpetuation of the species, since its currently endangered. The aim of the study was to characterize

cambucazeiro fruits as the physical and physico-chemical properties and the correlation between these attributes.

2. MATERIALS AND METHODS

Seventy-five fruits were collected from the median part of cambucazeiro plants, of the native Mata Atlântica collection of the Department of Plant Science of the Federal University of Viçosa (UFV), in Viçosa, Minas Gerais, Brazil, at 20°45'S, 42°51'O and 649 m of altitude. Fruits were sampled in February 2015, when they reached full physiological maturation, i.e. with yellow-orange peel. According to Koppen classification, the climate of the region is of the Cwa type, tropical altitude with dry winters and hot and rainy summers [12]. The average rainfall of the last 30 years was 1,221 mm, average air temperature of 19°C and average humidity of 80%.

Right after collection, all fruits were taken to a post-harvest laboratory, where they were washed and sanitized. Then, fruits were characterized as the following physical attributes: longitudinal and transverse diameter, measured with a digital pachymeter; longitudinal/transverse ratio; total mass of pulp (epicarp + mesocarp + endocarp) and seed (g), weighted in a semi-analytical balance; pulp and seed percentage (%), obtained by the ratio between each part's mass by total fruit mass; pulp coloration (chromaticity, hue° angle and luminosity), determined by reflectometry utilizing the reflectometer Minolta (Color Reader CT-10).

The analyzed chemical properties were: soluble solids (°Brix), obtained by means of the direct reading of a digital refractometer; titratable acidity (mg of citric acid per 100 mL⁻¹ of juice), determined by the titration of 10 mL of homogenized juice with 90 mL of distilled water, using a 0.1N NaOH solution as titrant and adding three drops of phenolphthalein as an

indicator to the sample; soluble solids/acidity ratio; ascorbic acid content (mg of ascorbic acid in 100 g of pulp), determined by titration with Tillman reagent (2.6 dichlorophenolindophenol - sodic salt at 0.1%) [13]; and the total carotenoids content (mg 100 g⁻¹ of pulp), determined through the extraction of approximately 2 g of mashed pulp with the addition of 50 mL of acetone 80 % and kept for 48 h inside a refrigerator at 4°C. Subsequently, the absorbance of the obtained extract was read in a spectrophotometer at the wavelength of 470.0 646.8 and 663.2 nm, and the carotenoids levels were determined [14].

Means were determined for all evaluated characteristics, as well as minimum and maximum values, standard deviation and variation coefficient. Then, all data underwent a Pearson's correlation analysis, among all variables. All statistical analysis was performed with the statistical program "GENES" [15].

3. RESULTS AND DISCUSSION

Average values of cambucá physical characteristics are presented in Table 1. Longitudinal and transverse fruits' diameter were 37.76 and 44.36 mm and the variation coefficient of 4.58 and 4.14% (considered low for these attributes), respectively, with the ratio longitudinal/transverse varying from 0.82 to 0.87, with average value of 0.85. These results indicate that this fruit has a flattened shape, due to a lower longitudinal diameter compared to the transverse, resulting in a ratio of these characteristics lower than 1. In "cagaita" (*Eugenia dysenterica* DC.), a species of the same family of the cambucá was observed lower average values of longitudinal (34.7 mm) and transverse (31.6 mm) diameter [10].

Fruits mass ranged from 35.21 to 55.42 g, with an average of 44.12 g, with a great homogeneity in evaluated data (VC = 13.24% and $\sigma = 5.88$ g). For *Psidium friedrichsthalianum* an average mass observed was of 42.19 g, ranging from 18.02 to 83.82 g [16], while in *Campomanesia phaea*, the fruit mass ranging from 21.2 to 82.33 g, with an average of 42.19 g [8]. These results indicate that cambucazeiro fruits' average mass is similar to other species of the Myrtaceae family.

Pulp mass presented a variation from 28.92 to 45.34 g, with an average of 36.46 g, while seed mass was 7.94 g, ranging from 15.66 to 20.41 g (Table 1). Given these results, the higher pulp

mass positively influenced on the increase in pulp percentage of fruits (Table 1). Pulp percentage was 82.15%, while seed was 17.84%. These values suggests that cambucazeiro fruits are fleshy, which is very interesting both for the industry and in natura consumption. Similar values to our results are observed in other Myrtaceae species, as in cagaita fruits, with pulp yield of 86.43% [10] and, in jabuticabeira fruits with 62.54% of the total fruits' mass corresponding to the pulp [3].

Color characterization variables such as chromaticity, hue angle and pulp luminosity are presented in Table 2. Mean values were 15.35 ($\sigma = 2.76$ and VC = 18.01%); 76.68 ($\sigma = 3.15$ and VC = 4.11) e 43,67 ($\sigma = 2.38$ and VC = 18.01%), with low coefficients of variation. Based on these variables, cambucazeiro fruits were characterized with yellow-orange coloration, in both peel and pulp. In the literature the cambucá fruit is described as yellow-orange, with little intensity of color and brightness [1].

In native and exotic Amazon fruits was observed varying colors from yellow with slightly green to yellow with little orange (e.g. noni, murici, abiu, yellow mombin, cajarana, cashew, bacuri, cupuacu and soursop), with hue angle close to 70, indicating yellow shades [17].

Chemical properties of the cambucá fruits are presented in Table 3. Soluble solids variation ranged from 8.72 to 12.60, with an average of 10.53 °Brix, while the titratable acidity contents varied from 0.71 to 2.07%, with 1.34% average of citric acid. Soluble solids content ranging from 7.3 to 13.30° Brix and titratable acidity of 0.65 to 3.48% of citric acid are reported in the literature [8]. Due to the acidic character of the cambucá fruits, there is a possibility of use on fruits processing, such as concentrated juices and fermented beverages [18].

The average content of ascorbic acid in cambucá fruits was 7.84%, while the carotenoids was 0.21 mg.100 g⁻¹ of pulp. Ascorbic acid and carotenoids are important for preventing and reducing lipids, proteins and nucleic acids' oxidative damage by reactive oxygen species, where free radicals are found [19].

Despite presenting antioxidative substance, determined by the contents of ascorbic acid and carotenoids, these values are below the observed for other native fruit species in Brazil, such as the mangaba (*Hancornia speciosa*), jabuticaba, caju (*Anacardium occidentale*) and

umbu (*Phytolacca dioica*), all of which presents carotenoids varying from 0.3 to 1.0 mg.100g⁻¹ of pulp and ascorbic acid contents of 18.4 to 23.8% [11]. These contents are also below the levels found in traditional fruits, such as mango, banana, papaya and guava, being the last one of the same family as cambucá [20,21,22,23].

Averages for the physical and physico-chemical characteristics of Myrtaceae species *Plinia cauliflora* (Jabuticaba), *Camponesia phaea* (Cambuci), *Eugenia dysentery* Dc. (Cagaita), *Psidium friedrichsthalianum* and *Eugenia uniflora* described in the literature [3,8,10,17,24] and the values verified in the present study are summarized in Table 4. There is quite diversity of sizes, forms, colors, organic acids and antioxidant compounds in species of Myrtaceae. Attractive coloration, the yield of pulp superior to

80%, fresh mass superior to the other species and characteristics of content of soluble solids, titratable total acidity similar to those verified for species already commercialized as the Jabuticaba and the cagaita indicate potential of commercial exploitation fruit of Cambucá.

Pearson's correlation values for physical and physico-chemical characteristics of cambucá fruits are presented in Table 5. It was verified that the total mass has high positive correlation with the following variables: longitudinal diameter (0.94), transverse diameter (0.94), pulp mass (0.99) and seed mass (0.85). On the other hand, negative correlations were verified with titratable acidity (0.71) at 1% significance, fruits with higher mass presented lower citric acid contents, represented by titratable acidity, contributing to the increase in the ratio SS/TA (0.66).

Table 1. Average values with respective minimum and maximum values, standard deviation (σ) and variation coefficient (VC) of the characteristics: longitudinal and transverse diameter, longitudinal/transverse ratio, total mass, pulp mass, seed mass, pulp and seed percentage of cambucá fruits (*Plinia edulis*)

Characteristics	Mean	Minimum	Maximum	Σ	VC (%)
Longitudinal diameter (LD) (mm)	37.76	34.74	40.91	1.73	4.58
Transverse diameter (TD) (mm)	44.36	41.26	47.79	1.93	4.14
Ratio LD/TD	0.85	0.82	0.87	0.20	2.31
Total mass (g)	44.12	35.21	55.42	5.88	13.24
Pulp mass (g)	36.46	28.92	45.34	4.73	12.97
Seed mass (g)	7.94	6.18	11.31	1.43	18.03
Pulp percentage (%)	82.15	79.58	84.33	1.65	9.24
Seed percentage (%)	17.84	15.66	20.41	1.65	2.01

σ : standard deviation, VC: variation coefficient.

Table 2. Average values with respective minimum and maximum values, standard deviation (σ) and variation coefficient (VC) of the characteristics: chromaticity, hue^o angle (HUE) and luminosity of the pulp of cambucá fruits (*Plinia edulis*)

Characteristics	Mean	Minimum	Maximum	Σ	VC (%)
Chromaticity	15.35	11.36	20.06	2.76	18.01
HUE	76.68	72.50	83.03	3.15	4.11
Luminosity	43.67	39.26	47.60	2.38	18.01

σ : standard deviation, CV: variation coefficient

Table 3. Average values with respective minimum and maximum values, standard deviation (σ) and variation coefficient (VC) of the characteristics: soluble solids, titratable acidity, soluble solids/titratable acidity ratio, ascorbic acid and carotenoid levels of the pulp of cambucá fruits (*Plinia edulis*)

Characteristics	Mean	Minimum	Maximum	σ	VC (%)
Soluble solids (°Brix)	10.53	8.72	12.60	1.00	9.54
Titratable acidity (% of citric acid)	1.34	0.71	2.07	0.48	35.58
Soluble solids/titratable acidity ratio	8.82	4.74	13.22	3.12	35.35
Ascorbic acid levels (% of ascorbic acid)	7.84	6.16	9.91	1.22	15.63
Carotenoid (mg 100 g ⁻¹ of pulp)	0.21	0.13	0.29	0.04	21.72

σ : standard deviation, CV: variation coefficient

Table 4. Average values of the characteristics: longitudinal diameter (LD), transverse diameter (TD), longitudinal/transverse diameter ratio (DL/DT), total mass (TM), pulp mass (PM), seed mass (SM), pulp percentage (PP), Seed percentage (SP) soluble solids (SS), titratable acidity (TA), titratable acidity (TA), soluble solids/titratable acidity ratio (SS/TA), ascorbic acid content (AA), carotenoids (CT), chromaticity (CRO), hue° angle (HUE) and luminosity (LUM) of some species Myrtaceae

Characteristics	Specie of Myrtaceae						
	Cambucá (<i>Plinia edulis</i>)	Jabuticaba (<i>Plinia cauliflora</i>) [3]	Cambuci (<i>Camponesia phaea</i>) [8]	Cagaita (<i>Eugenia dysenterica</i> DC.) [9]	Cagaita (<i>Eugenia dysenterica</i> DC.) [10]	Goiaba-da-Costa- Rica (<i>Psidium friedrichsthali anum</i>) [17]	Pitanga (<i>Eugenia uniflora</i> L.) [24]
LD (mm)	37.76	18.25	35.92	27.36	34.7	35.68	14.18
TD (mm)	44.36	18.18	50.10	32.53	31.6	42.9	19.27
DL/DT	0.85	1		0.84	1.1	0.83	0.73
TM (g)	44.12	3.86	42.25	18.08	25.11	42.16	2.79
PM (g)	36.46	2.41		12.48	21.72	39.72	2.21
SM (g)	7.94	0.33		2.8	3.39	2.47	0.58
PP (%)	82.15	62.54		69.03	86.43	94.21	79.46
SP (%)	17.84	8.66		15.49	13.5	5.86	20.54
SS (°Brix)	10.53		10.38	7.18			10.88
TA (% of citric acid)	1.34		2.71	1.48			1.87
SS/TA	8.82		4.75	4.85			5.85
AA (% of ascorbic acid)	7.84		78.95		34.11		
CT (mg 100 g ⁻¹ of pulp)	0.21				0.77		
CRO	15.35			49.65			
HUE	76.68			33.25			
LUM	43.67			82.63			

Table 5. Estimates of Pearson's correlation coefficients between the characteristics: total mass (TM), longitudinal diameter (LD), transverse diameter (TD), pulp mass (PM), seed mass (SM), pulp yield (PY), longitudinal/transverse diameter ratio (LD/TD), soluble solids (SS), chromaticity (CRO), luminosity (LUM), hue° angle (HUE), titratable acidity (TA), ascorbic acid content (AA), carotenoids (CT) and soluble solids/titratable acidity ratio (SS/TA) of cambucá fruits (*Plinia edulis*)

	LD	TD	PM	SC	PY	LD/TD	SS	CRO	LUM	HUE	TA	AA	CT	SS/TA
TM	0.94 **	0.94 **	0.99 **	0.85 **	-0.22	0.17	-0.32	-0.63 *	-0.48	0.41	-0.71 **	0.33	-0.29	0.66 **
LD		0.87 **	0.91 **	0.86 **	-0.32	0.44	-0.31	-0.68 **	-0.45	0.34	-0.72 **	0.47	-0.41	0.80 **
TD			0.92 **	0.83 **	-0.26	-0.07	-0.33	-0.50 **	-0.30	0.33	-0.52 *	0.19	-0.21	0.53 *
PM				0.75 **	-0.06	0.15	-0.26	-0.64 **	-0.53 *	0.38	-0.68 **	0.35	-0.28	0.62 *
SM					-0.70 **	0.22	-0.43	-0.46	-0.24	0.44	-0.64 **	0.20	-0.26	0.65 **
PY						-0.18	0.39	0.00	-0.20	-0.24	0.25	0.07	0.04	-0.34
LD/TD							-0.03	-0.46	-0.36	0.07	-0.51 *	0.59 *	-0.45	0.64 *
SS								-0.27	-0.04	0.05	0.16	0.52 *	-0.10	-0.50
CRO									0.53 *	-0.47	0.77 **	-0.82 **	0.48	-0.56 *
LUM										-0.14	0.47	-0.49	0.32	-0.43
HUE											-0.52 *	0.14	-0.05	0.11
TA												-0.44	0.32	-0.65 **
AA													-0.55 *	0.48
CT														-0.46

The superscript indexes *, ** and ns indicate correlation coefficients significant at 5 %, at 1 % and non-significant respectively

This results of positive correlation between total mass, longitudinal and transverse diameter, seed mass, and mass pulp, were similar of the ones verified for cagaita fruits [12]. Para *Psidium friedrichsthalianum* also verified high positive correlation between fruit mass and physical properties of the [16]. These correlations are important, since they indicate that plants selection with fruits of higher mass may be performed from the measurement of the diameter still in the field, without the need of weighting the fruits [10].

The variable chromaticity correlated negatively with most of the physical variables, beyond hue angle, vitamin C and ratio AT/SS of cambucá fruits, and positively with (although low) luminosity (-0.53) and titratable acidity (-0.77). The hue angle associated only with the acidity, demonstrating that the higher the pulp's angle, the lower is the acidity. The variables chromaticity, luminosity and hue° angle are used to describe the color of fruits, which directly influences consumer's choice.

Titratable acidity negatively correlated with the observed physical characteristics, except for pulp yield, which was not influenced by this variable. It was also observed a positive correlation between ascorbic acid content and the chemical variables, soluble solids and carotenoids. These results indicate that fruits with higher levels of ascorbic acid tend to present increased soluble solids and carotenoids. The ratio soluble solids/titratable acidity was positively correlated with total mass, longitudinal and transverse diameter, pulp and seed mass, while negatively with titratable acidity.

Positive correlations between soluble solids, ascorbic acid levels and carotenoids and the ratio soluble solids/titratable acidity was observed in papaya fruits. Significant associations of these variables are interesting because they allow genotype selection with greater levels of soluble solids and carotenoids [22].

The results obtained in fruits' characterization studies provide the establishment of important techniques of production for fruits, allowing the implantation of technified native fruit trees, where a difficulty of crops establishment exists, due to lack of information [16]. Thus, scientific information is important to establish cambucazeiro crops, enabling commercial

exploitation of its fruits and reducing the extinction risk of this species.

4. CONCLUSION

Cambucá fruits, a native species of the Brazilian Atlantic forest, present physical and physical-chemical characteristics that allow the commercial exploitation in the national market as well as internationally, being able to be commercialized as exotic fruits. The characteristics evaluated are within the standards observed for other native species already exploited commercially. Positive correlations verified for most of the characteristics indicate the possibility of genetic improvement of the species through the selection of more productive genotypes, without prejudice to the characteristics of the fruit flavor.

ACKNOWLEDGEMENTS

The authors thank Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brazil (CAPES) - Finance Code 001 for the financing of the research.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Lorenzi H, Bacher L, Lacerda M, Sartori S. Frutas brasileiras e exóticas cultivadas. 1st Ed. São Paulo: Instituto Plantarum de Estudos da Flora; 2006.
2. Danner MA, Citadin I, Sasso SAZ, Sachet MR, Ambrósio R. Phenology of blooming and fruiting of *Myrtaceae* native species of araucaria forest. Rev Bras Frut. 2010;32(1):291-295. DOI: 10.1590/S0100-29452010005000008
3. Guedes MNS, Rufini JCM, Azevedo AM, Pinto NAVD. Fruit quality of jaboticaba progenies cultivated in a tropical climate of altitude. Fruits. 2014;69(6):449-458. DOI: 10.1051/fruits/2 014030
4. Ishikawa T, Kato ETM, Yoshida M, Kaneko TM. Morphoanatomic aspects and phytochemical screening of *Plinia edulis* (Vell.) Sobral (*Myrtaceae*). Rev Bras Ciênc Farmac. 2008;44:515-520. DOI: 10.1590/S1516-93322008000300023

5. Alegretti AL, Wagner Júnior A, Bortolini A, Hossel C, Zanela, J, Citadin I. Biofilms and vacuum package in the storage of *Eugenia involucrata* DC. seeds. Rev Ceres. 2015;62(1):124-127.
DOI: 10.1590/0034-737X201562010016
6. Rocha WS, Lopes RM, Silva DB, Vieira RF, Silva JP, Agostini-Costa TS. Compostos fenólicos totais e taninos condensados em frutas nativas do cerrado. Rev Bras Frut. 2011;33(4):1215-1221.
DOI: 10.1590/S0100-29452011000400021
7. Hossel C, Oliveira JSMA, Wagner Júnior A, Mazaro SM, Citadin I. Cultural practice of root pruning during the transplante of native fruit plantlets. Rev Bras Frut. 2014;36(3):761-765.
DOI: 10.1590/0100-2945-278/13
8. Bianchini FG, Balbi RV, Pio R, Silva DF, Pasqual M, Vilas Boas EVB. Morphological and chemical characterization of the fruits of cambuci fruit tree. Bragantia. 2016;75(1):10-18.
DOI: 10.1590/1678-4499.096
9. Camilo YMV, Souza ERB, Vera R, Naves RV. Fruit characterization and progeny selection of cagaita (*Eugenia dysenterica* DC.). Científica. 2014;42(1):1-10.
DOI: 10.15361/1984-5529.2014v42n1p1-10
10. Cardoso LM, Martino HSD, Moreira AVB, Ribeiro SMR, Pinheiro-Sant'ana HM. Cagaita (*Eugenia dysenterica* DC.) of the Cerrado of Minas Gerais, Brazil: Physical and chemical characterization, carotenoids and vitamins. Food Research International. 2011;44(7):2151-2154.
DOI: 10.1016/j.foodres.2011.03.005
11. Rufino MSM, Alves RE, Brito ES, Pérez-Jiménez J, Saura-Calixto F, Mancini-Filho J. Bioactive compounds and antioxidant capacities of 18 non-traditional tropical fruits from Brazil. Food Chemistry. 2010;121:996-1002.
DOI: 10.1016/j.foodchem.2010.01.037
12. Alvares CA, Stape JL, Sentelhas PC, Gonçalves JLM, Sparovek G. Köppen's climate classification map for Brazil. Meteorol. Z. 2013;22(6):711-728.
DOI: 10.1127/0941-2948/2013/0507
13. AOAC - Association of Official Analytical Chemists. Official methods of analysis of AOAC international. 16th Ed. Maryland: AOAC; 1997.
14. Lichtenthaler HK. Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. Methods in Enzym. 1987;148:349-382.
DOI: 10.1016/0076-6879(87)48036-1
15. Cruz CD. GENES - A software package for analysis in experimental statistics and quantitative genetics. Acta Scient Agro. 2013;35(3):271-276.
DOI: 10.4025/actasciagron.v35i3.21251
16. Rebouças ER, Gentil DFO, Ferreira SAN. Physical characterization of fruits and seeds of Costa Rican Guavas Produced in Manaus, Amazonas. Rev Bras Frut. 2008;30(2):546-548.
DOI: 10.1590/S0100-29452008000200048
17. Canuto GAB, Xavier AAO, Neves LC, Benassi MT. Physical and chemical characterization of fruit pulps from Amazonia and their correlation to free radical scavenger activity. Rev Bras Frut. 2010;32(4):1196-1205.
DOI: 10.1590/S0100-29452010005000122
18. Chitarra MIF, Chitarra AB. Pós-colheita de Frutas e hortaliças: Fisiologia e manuseio. 2nd Ed. Lavras: UFLA; 2005.
19. Couto MAL, Canniatti-Brazaca SG. Quantification of vitamin C and antioxidant capacity of citrus varieties. Ciênc Tecnol Aliment. 2010;30(1):15-19.
DOI: 10.1590/S0101-20612010000500003
20. Amorim EP, Cohen KO, Amorim VBO, Paes NS, Sousa HN, Santos-Serejo JÁ, Silva SO. Characterization of banana accessions with base on functional compounds. Ciência Rural. 2011;41(4):592-598.
DOI: 10.1590/S0103-84782011005000042
21. Mendonça RD, Ferreira KS, Souza LM, Marinho CS, Teixeira SL. Physical and chemical characteristics of 'Cortibel 1' and 'Cortibel 4' guavas stored in environmental conditions. Bragantia. 2007;66:685-692.
DOI: 10.1590/S0006-87052007000400019
22. Reis RC, Viana ES, Jesus JL, Dantas JLL, Lucena RS. Physicochemical characterization of new hybrids and inbred lines of papaya. Pesq Agropec Bras. 2015;50(3):210-217.
DOI: 10.1590/S0100-204X2015000300004

23. Silva DFP, Siqueira DL, Rocha A, Salomão LCC, Matias RGP, Struiving TB. Genetic diversity among cultivars of mango based on fruit quality traits. Rev Ceres. 2012;59(2):225-232.
DOI: 10.1590/S0034-737X2012000200011
24. Dias AB, Carvalho MAP, Dantas ACVL, Fonseca VJA. Variabilidade e caracterização de frutos de pitangueiras em municípios baianos. Rev Bras Frut. 2011;33(4):1169-1177.
DOI: 10.1590/S0100-29452011000400015

© 2019 Souto et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history47242>